



# HIV TESTING GUIDELINES

## FOR THE PROVINCE OF BRITISH COLUMBIA



Office of the  
Provincial Health Officer

## CONTENTS

Background	2
Summary of Recommendations	3
Contributors	3
Who should be offered an HIV test?	4
Special considerations for testing	6
When to test after a possible exposure to HIV	6
Nominal vs. Non-nominal reporting	6
Other testing options	6
Point of care testing	6
Consent and HIV testing	6
Managing results	7
Non-attendance for positive results	7
Initial management of positive HIV test	7
Resources for clinicians	8
References	8

## BACKGROUND

Today, HIV infection is a chronic manageable medical condition. Early diagnosis and treatment with antiretroviral therapy (ART) allows HIV infected patients to live long and productive lives, and reduces transmission of the virus.

In British Columbia in 2014, there were an estimated 12,000 people living with HIV. Each year 200-300 people ranging in age from 13 to 81 years old are diagnosed with HIV infection.<sup>1</sup>

Despite advances in treatment, there continues to be significant and avoidable morbidity and mortality occurring amongst HIV infected individuals, much of which is attributable to late diagnosis. In the last decade in Vancouver, over 60% of diagnoses occurred after patients should already be on treatment.<sup>2</sup> In BC up to 17% of patients have advanced disease at the time of diagnosis.<sup>3</sup> Data from the United Kingdom<sup>4</sup>, United States<sup>5</sup>, and Vancouver<sup>6</sup> indicate that people diagnosed late in their infection have had multiple missed opportunities for earlier diagnosis in acute, community, and primary care settings.

This delay in diagnosis has consequences for individuals as well as for the community. For individuals diagnosed late, there is an impaired response to ART, as well as overall increased morbidity and mortality. For the community, late diagnosis contributes to the spread of HIV. It is estimated that 54% of new HIV infections occur via transmission from individuals who are unaware of their HIV status<sup>7</sup>. However, people who are diagnosed and are on effective treatment are significantly less likely to transmit infection to their partners.

To decrease late diagnoses, many jurisdictions, such as the United States<sup>8,9</sup>, the United Kingdom<sup>10</sup>, and France<sup>11</sup>, now recommend routine HIV testing in acute and primary care – in addition to existing approaches to HIV diagnosis. This approach recognizes that HIV testing based on the recognition of an individual's risk factors is insufficient to meet the goal of early diagnosis for all patients. It has been demonstrated to be highly acceptable to patients and health care providers, and effective in reaching and diagnosing patients who were not benefiting from early diagnosis and treatment.<sup>12</sup>

Health care providers may not perceive a risk or understand an individual to be at risk for HIV, and individuals may not disclose their reasons for testing or their risks for HIV. Requiring patients to disclose a risk to receive an HIV test can be an unintended barrier to testing and a missed opportunity for diagnosis. As such, an HIV test should be ordered whenever an individual requests it.

Experience with antenatal screening has shown that routine testing of a large group of individuals based on demographic factors (and not on risk factors), is considered generally acceptable and is a critical element in effective prevention of transmission of HIV.

These guidelines articulate current HIV testing recommendations for British Columbia. Each component of these guidelines will be evaluated with ongoing monitoring and assessment. As with all guidelines, HIV testing recommendations will evolve over time.

We recommend that health care providers know the HIV status of all patients under their care.

## Specifically, we recommend that providers offer an HIV test

- **Routinely**, every five years, to all patients aged 18-70 years
- **Routinely**, every year, to all patients aged 18-70 years who belong to populations with a higher burden of HIV infection
- **Once** for patients older than 70 years of age, if HIV status is not known

**AND** offer an HIV test to patients including adults 18-70, youth and the elderly, whenever

- Ordering diagnostic bloodwork for a new or worsening medical condition
- They present with symptoms of HIV infection or advanced HIV disease
- They or their providers identify a risk for HIV acquisition
- They request an HIV test
- They are pregnant
- You test for or diagnose a sexually transmitted infection (STI), hepatitis C, hepatitis B or tuberculosis

## CONTRIBUTORS

Dr. Perry Kendall, Provincial Health Officer (Guideline Sponsor)  
 Dr. Réka Gustafson, Vancouver Coastal Health (VCH) (Co-Chair)  
 Dr. Gina Ogilvie, BC Center for Disease Control (BCCDC) (Co-Chair)  
 Dr. Evan Adams, Deputy Provincial Health Officer (Public Health)  
 Jennifer Beaveridge, VCH (Nurse Practitioner)  
 Dr. Nora Cummins (Internal Medicine)  
 Dr. Charmaine Enns, Island Health (Public Health)  
 Dr. Mark Gilbert, BCCDC (Provincial Epidemiology)  
 Dr. David Hall, VCH (Family Practitioner, HIV treatment expert)  
 Dr. Oona Hayes, Island Health (Family Practitioner)  
 Dr. Pamela Kibsey, Island Health (Laboratories)

Dr. Kristen Korol, Island Health (Sexual Health Expert)  
 Dr. Mel Krajden, Provincial Health Services Authority (Laboratories)  
 Dr. Sandra Lee, Guidelines and Protocols Advisory Committee (Family Practitioner)  
 Dr. Richard Lester, BCCDC (STI/HIV Expert)  
 Dr. Gregory Linton, Northern Health (NH) (Family Practitioner)  
 Dr. Heather McDonald, Interior Health (IH) (Family Practitioner)  
 Kendra McPherson, VCH (Project Manager)  
 Dr. Val Montessori, BC Center for Excellence in HIV/AIDS (HIV Treatment Expert)  
 Dr. Michael Murphy, IH (Family Practitioner)

Dr. Fraser Norrie, VCH (Expert in care of MSM)  
 Ciro Panessa, Ministry of Health (Public Health)  
 Dr. Gurdeep Parhar, Fraser Health (FHA) (Family Practitioner)  
 Amrit Rai, FHA (Nursing)  
 Manik Saini, MoH (Communicable Disease Prevention)  
 Margot Smythe, VCH (Nursing)  
 Dr. Isaac Sobol, First Nations Health Authority (FNHA) (Aboriginal Health)  
 Denise Thomas, FNHA (Aboriginal Health)  
 Hannah Varto, VCH (Nurse Practitioner)  
 Dr. Barbra Arnold, BCCDC (Writer and Editor)

## ROUTINE TESTING FOR HIV - INDIVIDUALS PRESENTING FOR CARE

### RECOMMENDATION

Offer an HIV test to all individuals 18-70 years of age in your practice. For patients over 70 whose HIV status is not known, test once and then test if indicated by one of the considerations below.

### FREQUENCY

After an initial HIV test in all patients, repeat the HIV test at a frequency of every five years, or earlier if another indication for HIV testing is identified. The optimum frequency of testing in British Columbia's population is not yet determined, and the recommended frequency may change over time.

Some populations in BC are at increased vulnerability and experience a higher burden of HIV infection and morbidity.

Offer patients who are members of these populations HIV testing annually, or earlier if another indication for HIV testing is identified. These populations include:

- *Gay men*
- *People who inject drugs*
- *People who work in the sex trade*
- *People from endemic countries\**
- *Aboriginal people\*\**

### IN PRACTICE

Offer an HIV test when doing blood work for another reason

Offer as part of new patient intake

Offer when you do not have an HIV result for your patient in the past five years

\* IN 2014 COUNTRIES WHERE HIV IS ENDEMIC INCLUDE COUNTRIES OF THE CARIBBEAN AND SUB-SAHARAN AFRICA.

\*\* BC'S ABORIGINAL POPULATION, LIKE OTHER POPULATIONS WITH A HIGHER BURDEN OF DISEASE, IS DIVERSE AND HAS A RANGE OF HIV PREVALENCE. AS WITH OTHER POPULATIONS HAVING A HIGHER BURDEN, RECOMMENDATIONS ON TESTING FREQUENCY MAY BE SUBJECT TO CHANGE.

## CHANGE IN HEALTH STATUS

### RECOMMENDATION

Offer an HIV test to all patients, including those over 70 and youth, whenever ordering diagnostic bloodwork for a new or worsening medical condition.

### FREQUENCY

As clinically indicated

### IN PRACTICE

As HIV can have an array of nonspecific presentations, include HIV infection in the differential diagnosis for all patients, whether or not an individual risk for HIV acquisition has been identified. This includes when

a) a patient presents with symptoms that warrant laboratory investigation. Such conditions include **but are not limited to**; fever of unknown origin, mononucleosis-like syndrome, pneumonia, unexplained weight loss, unexplained hematological abnormality, and fatigue or failure to thrive.

b) a patient presents with symptoms associated with HIV infection or immune compromise. These symptoms include, but are not limited to lymphadenopathy, herpes zoster, recurrent and/or chronic herpes simplex infection, anogenital warts, anal cancer, cervical cancer, molluscum contagiosum, unexplained or recalcitrant prolonged diarrhea, unexplained peripheral neuropathy, Bell's palsy, oral candidiasis, oral hairy leukoplakia, seborrheic dermatitis, fungal infections, recurrent bacterial infections (e.g. cellulitis, folliculitis, pneumonia, bronchitis), unexplained dementia, aseptic meningitis, B-cell lymphoma, Kaposi's sarcoma or opportunistic infection indicative of immunodeficiency.

## INDIVIDUAL RISK TRIGGERED TESTING

### RECOMMENDATION

Offer an HIV test to all patients, including youth and people over 70 years of age, when a risk for HIV infection is identified

Offer an HIV test every time you test for or diagnose:

- *A sexually transmitted infection*
- *Hepatitis C*
- *Hepatitis B*
- *Tuberculosis*

### FREQUENCY

As clinically indicated by an identified risk

If an ongoing risk is present, test every 3-6 months

After an incarceration

If a recent high-risk exposure has occurred, or acute HIV infection is suspected, repeat at 4 and 12 weeks

### IN PRACTICE

If a recent high-risk exposure has occurred, or acute HIV infection (seroconversion) is suspected, indicate “query acute HIV” on the test requisition

---

## PATIENT INITIATED TESTING

### RECOMMENDATION

Order an HIV test whenever a patient requests it

### FREQUENCY

Whenever a patient asks

### IN PRACTICE

Individuals may not disclose their reasons (or their risks) for testing for HIV. Eliciting these reasons or risks may be a barrier to testing

---

## ANTENATAL SCREENING FOR HIV

### RECOMMENDATION

Offer an HIV test as part of routine antenatal care for all patients

### FREQUENCY

Once for every pregnancy

*If at ongoing risk, repeat in the third trimester*

## CONSENT & HIV TESTING

The purpose of a pre-test discussion is to establish informed consent. For HIV testing, obtaining informed consent is the same as for any other diagnostic test or treatment. As with other diagnostic tests, if the pretest probability of a positive result is high, more extensive discussion may be warranted. See: BCCDC Communicable Disease Control Manual Chapter 5 - Section II - HIV/AIDS. If a patient declines an HIV test, the reason for refusing the test should be explored to ensure it is not due to false information about HIV infection or the consequences of testing.

## SPECIAL CONSIDERATIONS FOR TESTING

### When to test after a possible exposure to HIV<sup>14</sup>

It is important to understand HIV testing window periods when considering an HIV test after a possible exposure. Traditionally, waiting 3 months after exposure was recommended; however, newer 4th generation HIV tests have shortened the window period. The average window period for 4th generation enzyme immunoassays (EIA) tests (which detect p24 antigen and HIV antibodies) is 16-18 days.

Diagnostic methods and technologies continue to evolve and the window period may be shortened in the future.

Most patients can be tested at 4 weeks following exposure (>95% of infected individuals will have detectable antibodies at this time). If negative, repeat testing at 3 months is recommended (>99% of infected individuals will have a positive EIA at this time).

If a recent high-risk exposure has occurred, or acute HIV infection (seroconversion) is suspected, test now. Please indicate “query acute HIV” on the laboratory requisition. Molecular detection methods can be added to standard serology.

Results of these tests require interpretation within the clinical and epidemiological context of the patient. A negative result in an individual with a high likelihood of being HIV positive can be reviewed with the medical microbiologist responsible for the testing laboratory.

### Non-nominal Reporting

Non-nominal reporting is available in British Columbia, meaning that the full name of the individual is not reported to public health. The first name, initials, and date of birth are reported to the Medical Health Officer for surveillance purposes. The full name of the individual is still on the specimen sent to the laboratory.

### Other Testing Options

Occasionally, providers may encounter a patient who has concerns about confidentiality in HIV testing.

Testing using an alias or initials only is available at certain sites in British Columbia. See: <http://smartsexresource.com/get-tested/hiv-testing>

A pilot of anonymous testing is underway in British Columbia. With an anonymous HIV test, the test is identified by a code known only to the patient. No identifiable or contact information is collected and the person being tested must provide his or her anonymous testing code in order to receive the result.

Patients seeking an anonymous HIV test can access this test by: <http://smartsexresource.com/topics/hiv-anonymous-testing>.

These forms of testing are not ideal from a public health perspective. However, if confidentiality is a barrier to testing, these options are available.

### Point of Care Testing

In British Columbia, point of care testing is used in the outreach setting. It has a similar sensitivity to traditional 3rd generation HIV testing. Confirmatory testing is required for indeterminate, invalid, and preliminary positive results. Point of care testing is insufficient to rule out acute HIV infection and is not recommended for those with a recent exposure.

For more information on POC testing, please see:

<http://www.bccdc.ca/SexualHealth/Programs/ProvincialPointofCareHIVTestingProgram/default.htm>

## MANAGING RESULTS

### FOR INDIVIDUALS WHO TEST HIV NEGATIVE

A separate post-test visit is not necessary. Results can be handled as any other negative result is handled in your office. Giving a negative HIV test result can be an opportunity to educate about risk and risk reduction, if indicated.

### FOR INDIVIDUALS WHO TEST HIV POSITIVE

Given the availability of resources, medical education, and support, a primary care provider, who is willing to do so can provide the necessary care in the majority of situations. In a scenario where that is not possible, or in the advanced stages of infection, consideration should be given to shared patient management with clinicians experienced in HIV.

A positive result should be given face to face in a confidential environment and in a clear and direct manner, as is good clinical practice for any situation where bad news is being conveyed.

Consult with HIV specialty care, if required.

Provide linkages to support and care including medical, emotional, nutritional, psychosocial, spiritual and financial, as with other serious medical diagnoses.

As HIV is a reportable infection, a positive HIV result will be sent to local public health officials. Public health nurses have a wealth of information and resources and can be part of your clinical team in initial management of someone recently diagnosed with HIV infection. They will be involved in the care of individuals with a new HIV infection and care of their partners. For example, public health nurses can be involved with delivery of the diagnosis, partner notification, and linkage to care.

Counsel on risk reduction. Individuals with recently acquired HIV infection have a much higher potential for transmitting the virus, and safer sexual practices are essential during the acute phase of HIV infection.

With current laboratory standards, false positive tests are exceedingly rare. Nevertheless, since a potential for error exists for diagnostic systems, a second test should be performed to confirm the diagnosis.

### NON-ATTENDANCE FOR POSITIVE RESULTS

Notify Public Health as they will have the resources and experience to assist with this issue.

### INITIAL MANAGEMENT OF POSITIVE HIV TEST

Review the case with your local Public Health office for consultative expertise related to partner notification and if required, linkage to care and supports.

### PROVIDING CARE TO PATIENTS DIAGNOSED WITH HIV

Allow sufficient time to discuss the diagnosis with your patient.

Refer or consult with a clinician experienced in the treatment and management of HIV infection, if appropriate.

Order the standard baseline testing following HIV diagnosis as follows:

- Repeat HIV Antibody (HIV test)
- HIV plasma viral load
- CD4/CD8 cell counts and ratio
- CBC and differential
- ALT, AST, Alk Phos, GGT, LDH, Bilirubin, INR, and Amylase
- Creatinine (eGFR), Na, K, Cl, HCO<sub>3</sub>, BUN
- Urinalysis
- Syphilis screen (RPR)
- Urine NAT for Gonorrhea and Chlamydia
- Hepatitis A Total Antibody
- Hepatitis B (HBsAg, anti-HBs Ab, anti-HBc Ab Total)
- Hepatitis C Ab, Hepatitis C RNA
- Toxoplasma IgG
- Pregnancy test (if appropriate)

Offer ongoing support and assess the psychosocial impacts of a recent HIV diagnosis.

Discuss prevention of transmission and disclosure to past and potential future partners. In Canada, nondisclosure of a positive HIV status may have legal implications. These legalities are evolving. For more information, see:

<http://www.aidslaw.ca/site/sex-criminal-law-and-hiv-non-disclosure/>

Ongoing care by a primary care provider, with assistance as needed by an expert in HIV care, is extremely important in optimizing patient care.

Provide ongoing care guided by the primary care guidelines and therapeutic guidelines found at the British Columbia Centre for Excellence in HIV/AIDS website pages:

<http://www.cfenet.ubc.ca/therapeutic-guidelines/primary-care>

<http://www.cfenet.ubc.ca/therapeutic-guidelines/adult>

## RESOURCES FOR CLINICIANS

### Local Public Health

#### Fraser Health

Fraser East (Chilliwack) 604-864-3437  
(Abbotsford) 604-702-4921  
Fraser North 604-777-6709  
Fraser South 604-587-7902

#### Vancouver Coastal Health

VCH Communicable Disease Control 604-675-3900  
(Ask for Communicable Disease Nurse on call)

#### Interior Health

Communicable Disease Unit 1-866-778-7736

#### Island Health

Central Island 1-866-770-7798  
North Island 1-877-887-8835  
South Island 1-866-665-6626

#### Northern Health

Northwest (Ask for the Designated Nurse) 250-631-4228  
Northern Interior 778-349-2793  
Northeast (Ask for the Designated Nurse) 250-719-6500

### BC Centre for Disease Control STI/HIV Prevention and Control

604-707-5600

<http://www.bccdc.ca>

### British Columbia Centre for Excellence in HIV/AIDS (BC-CfE) Rapid Expert Advice and Consultation for HIV (REACH) Line

<http://www.cfenet.ubc.ca/REACH>

604-681-5748 (Vancouver)

1-800-665-7677 (Outside Vancouver)

### BC Women's Hospital & Health Centre Oak Tree Clinic

<http://www.bcwomens.ca/Services/HealthServices/OakTreeClinic/default.htm>

604-875-2212

1-888-711-3030 (Toll Free in BC)

## REFERENCES

1. BC Centre for Disease Control. "HIV in British Columbia: Annual Surveillance Report 2012." 2013. Accessed: May 4<sup>th</sup>, 2014 <<http://www.bccdc.ca/util/about/annreport/default.htm>>.
  2. Tim Chu, Ellen Demlow, Reka Gustafson, Jat Sandhu. "STOP HIV/AIDS Semi-Annual Monitoring Report." 2013. Accessed: May 4<sup>th</sup>, 2014 <[http://www.vch.ca/media/Revised\\_H1\\_Semi\\_Annual\\_Monitoring\\_Report\\_Through\\_June%20\\_30\\_2013\(1\).pdf](http://www.vch.ca/media/Revised_H1_Semi_Annual_Monitoring_Report_Through_June%20_30_2013(1).pdf)>.
  3. Claudia Rank, Elisa Lloyd-Smith, Mark Gilbert. "Advanced HIV Disease at the Time of Diagnosis in British Columbia 1995-2008." 2011. Accessed: May 4<sup>th</sup>, 2014 <[http://www.bccdc.ca/NR/rdonlyres/332498AE-1B79-4D29-8F3D-AD9034D1576F0/STI\\_HIVSpecialReport\\_20110401.pdf](http://www.bccdc.ca/NR/rdonlyres/332498AE-1B79-4D29-8F3D-AD9034D1576F0/STI_HIVSpecialReport_20110401.pdf)>.
  4. Ann K Sullivan, Hilary Curtis, Caroline A, Sabin, Margaret A Johnson. "Newly diagnosed HIV infections: review in UK and Ireland." *BMJ* 330 (2005): 1301.
  5. US Centers for Disease Control and Prevention. "Missed opportunities for earlier diagnosis of HIV infection-South Carolina 1997-2005." *MMWR* 55.47 (2006): 1269-72.
  6. Ellen Demlow, Lauren MacDonald, Afshan Nathoo, Tim Chu, Jat Sandhu, Reka Gustafson. "Missed Opportunities for HIV Diagnoses in Vancouver." *CAHR* 2013, Innovations to Address Complex Challenges. Vancouver: Canadian Journal of Infectious Diseases and Medical Microbiology, 2013. 37. Accessed May 5<sup>th</sup>, 2014 <[http://www.pulsus.com/journals/pdf\\_frameset.jsp?nlKy=3&atlKy=777&isArt=f&jnlAdvert=Infdis&adverif-HCTp=&supKy=516&sTitle=CAHR%20Abstracts%2C%20Pulsus%20Group%20Inc&VisitorType=>](http://www.pulsus.com/journals/pdf_frameset.jsp?nlKy=3&atlKy=777&isArt=f&jnlAdvert=Infdis&adverif-HCTp=&supKy=516&sTitle=CAHR%20Abstracts%2C%20Pulsus%20Group%20Inc&VisitorType=>)>.
  7. Marks G, Crepaz N, Janssen RS. "Estimating sexual transmission of HIV from persons aware and unaware that they are infected with the virus in the USA." *AIDS* (2006): 1447-1950.
  8. Bernard M. Branson, H. Hunter Handsfield, Margaret A. Lampe, Robert S. Janssen, Allan W. Taylor, Sheryl B. Lyss, Jill E. Clark. "Revised Recommendations for HIV Testing of Adults, Adolescents, and Pregnant Women in Health-Care Settings." *MMWR*. Vol. 55. RR-14. 22 September 2006. 1-17.
  9. US Preventive Services Task Force. "Screening for HIV: U.S. Preventive Services Task Force Recommendation Statement." April 2013. Accessed: May 5<sup>th</sup>, 2014 <<http://www.uspreventiveservicestaskforce.org/uspstf13/hiv/hivfinalrs.htm>>.
  10. British HIV Association, British Association of Sexual Health and HIV, British Infection Society. "UK National Guidelines for HIV Testing, 2008." September 2008. Accessed May 5<sup>th</sup>, 2014 <<http://www.bhiva.org/documents/guidelines/testing/glineshivtest08.pdf>>.
  11. Haute Autorité de Santé. "HIV Infection Screening in France-Screening Strategies." October 2009. Accessed May 5<sup>th</sup>, 2014 <[http://www.has-sante.fr/portail/upload/docs/application/pdf/2010-02/hiv\\_infection\\_screening\\_in\\_france\\_-\\_screening\\_strategies\\_-\\_executive\\_summary\\_2010-02-26\\_10-28-32\\_643.pdf](http://www.has-sante.fr/portail/upload/docs/application/pdf/2010-02/hiv_infection_screening_in_france_-_screening_strategies_-_executive_summary_2010-02-26_10-28-32_643.pdf)>.
  12. Health Protection Agency. "Time to test for HIV: Expanded healthcare and community HIV testing in England." 2010. Accessed May 5<sup>th</sup>, 2014 <<http://www.hpa.org.uk/Publications/InfectiousDiseases/HIVandSTIs/1011TimetotestHIVtesting/>>>.
  13. Reka Gustafson, Gina S Ogilvie, David Moore, Perry Kendall. "New HIV testing guidelines in BC." *BC Medical Journal* 54.4 (2014): 172-3.
  14. Mark Gilbert, Mel Krajden. "Don't wait to test for HIV." *BCMJ* 52.6 (2010): 308-9.
8. HIV TESTING GUIDELINES FOR THE PROVINCE OF BRITISH COLUMBIA