



BC Centre for Disease Control
AN AGENCY OF THE PROVINCIAL HEALTH SERVICES AUTHORITY

STI

Annual Report
2014

Contact Information

BC Centre for Disease Control
Clinical Prevention Services
655 West 12th Avenue
Vancouver BC V5Z 4R4
Phone: 604-707-5621
Fax: 606-707-5604
Email: CPSSurveillance@bccdc.ca

Date of publication: November 30, 2015
Report is available at www.bccdc.ca

Suggested citation:

BC Centre for Disease Control. (2015). STI in British Columbia: Annual Surveillance Report 2014.
Retrieved from <http://www.bccdc.ca/util/about/annreport/default.htm>



Table of Contents

Summary of Trends	4
Chlamydia	5
Genital Chlamydia by Region, Gender, and Age	5
Extra-genital Chlamydia	10
Perinatally-acquired Chlamydia	11
Lymphogranuloma Venereum	12
Gonorrhea	13
Genital Gonorrhea by Region, Gender, and Age	13
Extra-genital Gonorrhea	19
Perinatally-acquired Gonorrhea	19
Gonorrhea Antimicrobial Resistance	20
Pelvic Inflammatory Disease and Ectopic Pregnancy	22
Infectious Syphilis	24
Infectious Syphilis by Region, Gender, and Age	24
Infectious Syphilis by Ethnicity	29
Infectious Syphilis among Aboriginal Peoples	30
Infectious Syphilis by Exposure Category	31
Infectious Syphilis among Men who have Sex with Men	34
Stage of Infection at Time of Syphilis Diagnosis	36
Maternal and Early Congenital Syphilis	37
Endnotes	38
Contributors	40
Technical Appendix	41
Data Limitations	41
Case Definitions	42
Data Sources	43
Additional Notes	45

Summary of Trends

Genital Chlamydia

In 2014, the rate of genital chlamydia increased to 288.4 per 100,000 population, continuing the overall provincial trend of a steady increase since 1998.

- The highest rates were in Vancouver, Northwest, and Northeast Health Service Delivery Areas.
- Females continue to have higher rates of genital chlamydia infection compared to males.
- The highest rates were among young adults aged 20-29 years followed by adolescents aged 15-19 years.
- In 2014, there were 104 extra-genital infections identified and 1 perinatally-acquired infection.
- An increase in lymphogranuloma venereum (LGV) was identified in 2014. The number of LGV cases remains higher than historic levels. All cases were among men who have sex with men, many of whom are co-infected with HIV.

Genital Gonorrhoea

In 2014, the provincial rate of genital gonorrhoea increased to 38.9 per 100,000 population, continuing an overall steady increase since 1998.

- The highest rates were in Vancouver, Northwest, and Northern Interior Health Service Delivery Areas.
- Males have higher rates of infection compared to females and in 2014, rates in males increased while rates in females have been more stable.
- The highest rates of infection were among males aged 25-29 years and among females aged 20-24 years.
- In 2014, there were 362 extra-genital infections identified and no perinatally-acquired infection.
- Analysis of recent gonorrhoea antimicrobial resistance trends in BC demonstrates a reduction in the proportion of isolates with reduced susceptibility to cefixime and ceftriaxone since 2011.

Pelvic Inflammatory Disease and Ectopic Pregnancy

In 2013, the rate of physician billings and hospital admissions for women related to pelvic inflammatory disease was stable or decreased which is consistent with the overall decline in this potential complication of chlamydia or gonorrhoea infection over the past two decades. While hospitalizations of women related to ectopic pregnancy have shown a similar trend, the number of physician billings has shown a small but steady increase since 2003 in BC.

Infectious Syphilis

In 2014, the provincial rate of infectious syphilis decreased slightly to 11.9 per 100,000 population, continuing a steady increase since 2011.

- The highest rates of infection were in Vancouver, Fraser North, and North Shore/Coast Garibaldi Health Service Delivery Areas.
- In 2014, over 95% of cases were male, with highest rates observed in males aged 25-39 years. A slight increase in female cases was identified in 2014.
- The majority of cases in 2014 were among people identified as Caucasian (52%).
- Men who have sex with men (MSM) continue to comprise the greatest number of new infectious syphilis cases in BC (85% in 2014). Among MSM cases where HIV status is known, 60% were co-infected with HIV.

Chlamydia

Genital Chlamydia by Region, Gender, and Age


Genital chlamydia is the most commonly reported sexually transmitted infection in BC. As the majority of chlamydia infections are asymptomatic, the reported number of chlamydia infections is only a portion of the total population burden. If untreated, genital chlamydia may lead to complications such as pelvic inflammatory disease (a major cause of infertility, ectopic pregnancy, and chronic pain) in women and epididymo-orchitis in men.

Mirroring the national trend, genital chlamydia rates have steadily increased from 1998 to 2014 following a decline in the early 1990s (Figure 2). In 2014, the rate of genital chlamydia for BC increased to 288.4 (13,348 cases) from 269.6 (12,352 cases) per 100,000 population in 2013. The highest rates of genital chlamydia were in Northern and Vancouver Coastal Health Authorities (Figure 3). Rates among Health Service Delivery Areas varied with the highest rates in Vancouver and Northwest, and the lowest rates in Fraser East, East Kootenay and Fraser South (Figure 1).

Similar increases in chlamydia infections have been observed in high income countries around the world.^{1,2} There are multiple reasons for this increase, including increases in the sensitivity of laboratory tests and uptake of testing (e.g., the greater acceptability of urine-based tests among men) as well as provider screening practices. There may also be a paradoxical effect in which improvements in early screening and treatment for chlamydia over the past decades have resulted in individuals being less likely to develop full immunity thus consequently more susceptible to re-infection (known as the “arrested immunity” hypothesis^{3,4}). While data on population trends in sexual behaviour is not available for BC, it is possible that changes in behaviour, such as decreased condom use, may also be contributing to increasing chlamydia incidence.


Females continue to have approximately twice the diagnosis rate compared to males. The rate for females in 2014 increased to 353.0 (8,222 cases) from 339.7 (7,830 cases) per 100,000 population in 2013 (Figure 4). The greater number of infections among females is partially due to routine screening performed at the time of visits that were for other reasons (e.g., pap testing or contraception counselling). In 2014, the highest rates of chlamydia were among young adults aged 20-29 years followed by adolescents aged 15-19 years (Figure 6), driven primarily by the high rates of infection among young females. For females, the rate for the age group 15-19 years decreased slightly while the rates for all other age groups increased from 2013 to 2014 (Figure 7). Males aged 20-29 years had the highest chlamydia rates in 2014 compared with other age groups (Figure 8).

1. Genital chlamydia case reports in BC by health service delivery area, 2014




Rates calculated with population estimates released by BC Stats

2. Genital chlamydia case reports in BC and Canada, 1991 to 2014*




* 2014 Canadian rate is not available

3. Genital chlamydia case reports in BC by health authority, 2005 to 2014



4. Genital chlamydia case reports in BC by gender, 2005 to 2014




* Other - transgender and gender unknown

5. Genital chlamydia case reports in BC by age group and gender, 2014




* Other - transgender and gender unknown


6. Genital chlamydia case reports in BC by age group - total, 2005 to 2014



7. Genital chlamydia case reports in BC by age group - female, 2005 to 2014



8. Genital chlamydia case reports in BC by age group - male, 2005 to 2014



Extra-genital Chlamydia

In 2014, 104 extra-genital cases were identified (15 females, 89 males) which is an increase from 57 cases in 2013 (11 females and 46 males). As screening for chlamydia infections at extra-genital sites is not routine practice, these findings are strongly influenced by provider testing practices. Much of the increase observed since 2011 is likely due to increased awareness and testing for chlamydia in extra-genital sites. From 2005 to 2014, 347 infections were identified in specimens collected from the following sites: throat (222 cases, 64.0%), eye (94 cases, 27.1%), lung (1 case, 0.3%), and other sites (30 cases, 8.6%) (Table 9).

9. Extra-genital chlamydia case reports in BC by site/culture, 2005 to 2014

Gender	Site	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Female	Throat	0	1	2	0	0	0	0	6	5	9
	Eye	9	6	3	3	6	3	3	2	2	6
	Lung	0	0	0	0	0	0	0	0	0	0
	Other *	0	0	1	1	4	5	1	1	4	0
	Total	9	7	6	4	10	8	4	9	11	15
Male	Throat	4	4	1	6	2	6	1	52	42	80
	Eye	5	8	5	5	5	5	5	3	2	8
	Lung	0	0	1	0	0	0	0	0	0	0
	Other *	0	0	0	2	0	2	4	2	2	1
	Total	9	12	7	13	7	13	10	57	46	89
BC	Throat	4	5	3	6	2	6	1	59	47	89
	Eye	14	14	8	8	11	8	8	5	4	14
	Lung	0	0	1	0	0	0	0	0	0	0
	Other *	0	0	1	3	4	7	5	3	6	1
	Total	18	19	13	17	17	21	14	67	57	104


* Other - nasopharyngeal washing, lesion, fluid from groin mass lesion, pelvic fluid, and other

Total reports for BC is the sum of the following genders: female, male, transgender, and gender unknown

Perinatally-acquired Chlamydia

Genital chlamydia can be passed by pregnant women to their infants during delivery which can lead to ophthalmia neonatorum and pneumonia. One case of perinatally-acquired chlamydia infection was found in a conjunctival specimen in 2014. From 2005 to 2014, the majority of perinatal cases are from conjunctival specimens (17/18 cases, 94.4%) while one case (5.6%) was identified in a respiratory specimen (Figure 10).

10. Perinatally-acquired chlamydia case reports in BC by site, 2005 to 2014*



* 2014 conjunctival rate is not available

Lymphogranuloma Venereum

Lymphogranuloma venereum (LGV) is a sexually transmitted infection caused by *Chlamydia trachomatis* serovars L1, L2, and L3. The clinical presentation of LGV includes genital papules, ulcers, inguinal lymphadenopathy, and hemorrhagic proctitis. If left untreated, LGV can cause serious sequelae such as lymphatic obstruction or anogenital ulcerations. LGV can easily be misdiagnosed as other sexually transmitted infections or gastrointestinal disease.

LGV was first reported in Canada in 2003 and in BC in 2004. With increasing cases of LGV among gay, bisexual, and other men who have sex with men (MSM) in Europe and the US, provincial LGV surveillance commenced in 2004. Occurring in tandem with reports of increased transmission in the US⁵ and Europe⁶, an increase of LGV cases was observed in 2011 in BC, in part due to routine testing of rectal chlamydia specimens for LGV and augmented case-finding. In 2014, 26 LGV cases were identified, nine cases more than in 2013 (17 cases) (Figure 11).

From 2005 to 2014, 99 cases of LGV (82 confirmed, 17 probable) were reported in BC. Most cases (97 cases, 98.0%) were among MSM and diagnosed in Vancouver (86 cases, 86.9%). Of those with known HIV status, 69.9% (65/93 cases) were co-infected with HIV. Many cases (59/97 cases, 60.8%) presented with symptoms of proctitis. In 2014, the male rate of LGV in BC was 1.1 (26 cases) per 100,000 population and the average age was 44.0 years (range 20-68 years). The majority of cases in 2014 were among men who identified as Caucasian (19 cases, 73.1%).^{7,8}

11. Lymphogranuloma venereum case reports in BC, 2005 to 2014



Gonorrhoea

Genital Gonorrhoea by Region, Gender, and Age

Gonorrhoea infections may be asymptomatic or the symptoms may be mild, and as a result, the reports of diagnosed infections are only a portion of the total population burden. As with chlamydia, genital gonorrhoea if untreated can lead to pelvic inflammatory disease (and associated complications) in females. An individual infected with gonorrhoea is also at increased risk of acquiring HIV.^{9, 10}


Overall since 1998, the provincial genital gonorrhoea rate in BC has increased consistent with national rates (Figure 13). The genital gonorrhoea rate in BC increased to 38.9 (1,802 cases) in 2014 from 37.2 (1,704 cases) per 100,000 population in 2013. The highest rate was in the Vancouver Coastal Health Authority (Figure 14). Rates among Health Service Delivery Areas vary with the highest rates in Vancouver, Northwest and Northern Interior, and the lowest rates in East Kootenay, Central Vancouver Island and Kootenay Boundary (Figure 12). Reasons for the increase in gonorrhoea rates over time include the routine inclusion of gonorrhoea with chlamydia urine nucleic acid testing and increased acceptability of urine-based testing among men. It is also possible that changes in behaviour, such as decreased condom use, may be contributing to a true increase in incidence during this time period.

Males continue to have a rate two times greater than females (Figure 15). Since 2005, male gonorrhoea rates have been steadily increasing. In 2014, the rate among males increased to 56.4 (1,296 cases) from 51.7 (1,177 cases) per 100,000 population in 2013. Female gonorrhoea rates have gradually increased since 2005. The rate among females decreased slightly in 2014 to 21.6 (502 cases) from 22.6 (522 cases) per 100,000 population in 2013.


Similar to trends from 2005 to 2013, in 2014, the highest rates of gonorrhoea were among those aged 20-29 years (Figure 17). In 2014, the highest rates among males were in those aged 25-29 years (284 cases, 180.8 per 100,000 population) and among females in those aged 20-24 years (136 cases, 88.9 per 100,000 population) (Figure 16).

Gonorrhoea is more likely to be concentrated in sexually active networks and it is likely that the higher rates of gonorrhoea in males is, in part, due to higher rates of gonorrhoea among gay, bisexual, and other men who have sex with men (MSM). While provincial surveillance data does not permit identification of MSM cases, this has been observed in other jurisdictions.^{11, 12}


12. Genital gonorrhoea case reports in BC by health service delivery area, 2014



Southwestern BC Inset




Rate per 100,000 population by HSDA



ID	Health Service Delivery Area	Cases	Rate
11	East Kootenay	3	3.9
12	Kootenay Boundary	7	9.0
13	Okanagan	33	9.3
14	Thompson Cariboo Shuswap	57	25.8
21	Fraser East	73	25.3
22	Fraser North	208	32.4
23	Fraser South	264	34.0
31	Richmond	38	18.5
32	Vancouver	819	124.4
33	North Shore/Coast Garibaldi	65	23.0
41	South Vancouver Island	79	21.2
42	Central Vancouver Island	22	8.3
43	North Vancouver Island	15	12.4
51	Northwest	34	46.1
52	Northern Interior	61	43.0
53	Northeast	9	12.5


Rates calculated with population estimates released by BC Stats

13. Genital gonorrhoea case reports in BC and Canada, 1991 to 2014*




* 2014 Canadian rate is not available

14. Genital gonorrhoea case reports in BC by health authority, 2005 to 2014




15. Genital gonorrhoea case reports in BC by gender, 2005 to 2014



	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Reports - Female	275	340	429	537	503	434	459	449	522	502
Reports - Male	825	681	793	866	805	883	1,112	845	1,177	1,296
Reports - Other*	0	1	2	2	1	3	3	3	5	4
Rate - Female	13.0	15.9	19.8	24.5	22.7	19.3	20.3	19.6	22.6	21.6
Rate - Male	39.7	32.4	37.3	40.1	36.7	39.8	49.7	37.4	51.7	56.4

* Other - transgender and gender unknown


16. Genital gonorrhoea case reports in BC by age group and gender, 2014




	10 - 14 yrs	15 - 19 yrs	20 - 24 yrs	25 - 29 yrs	30 - 39 yrs	40 - 59 yrs	60+ yrs
Reports - Female	2	76	136	110	104	68	5
Reports - Male	0	49	235	284	346	334	47
Reports - Other*	0	0	1	1	2	0	0
Rate - Female	1.8	58.3	88.9	69.5	33.4	10.0	0.9
Rate - Male	0.0	34.7	142.5	180.8	113.2	50.4	9.1

* Other - transgender and gender unknown


17. Genital gonorrhoea case reports in BC by age group - total, 2005 to 2014



18. Genital gonorrhoea case reports in BC by age group - female, 2005 to 2014



19. Genital gonorrhoea case reports in BC by age group - male, 2005 to 2014



Extra-genital Gonorrhoea

In 2014, 362 cases were identified (20 females, 342 males) which was an increase from 219 cases (28 females, 188 males) in 2013. As screening for gonorrhoea infections at extra-genital sites is not routine practice, these findings are strongly influenced by provider testing practices, including increases in screening in the past few years. Of the 1,256 cases diagnosed from 2005 to 2014, cases were identified in the throat (1,204 cases, 95.9%), eye (26 cases, 2.1%), and other sites (18 cases, 1.4%). A small number of the diagnosed cases represented disseminated gonococcal infection (8 cases, 0.6%) (Table 20).

20. Extra-genital gonorrhoea case reports in BC by site/culture, 2005 to 2014

Gender	Site	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Female	Throat	14	16	15	3	7	8	12	8	26	19
	Eye	0	0	1	1	0	1	0	0	1	1
	Other *	0	0	4	0	0	1	0	0	0	0
	DGI **	2	0	0	1	0	0	0	1	1	0
	Total	16	16	20	5	7	10	12	9	28	20
Male	Throat	74	41	46	41	43	55	95	156	184	335
	Eye	1	0	1	1	5	4	0	1	2	6
	Other *	1	1	0	0	1	3	3	1	2	1
	DGI **	0	0	0	1	1	0	0	1	0	0
	Total	76	42	47	43	50	62	98	159	188	342
BC	Throat	88	57	61	44	52	63	107	165	213	354
	Eye	1	0	2	2	5	5	0	1	3	7
	Other *	1	1	4	0	1	4	3	1	2	1
	DGI **	2	0	0	2	1	0	0	2	1	0
	Total	92	58	67	48	59	72	110	169	219	362

* Other - superficial wound, sternoclavicular synovium, synovium joint fluid, abscess, blood, and elbow

** DGI - disseminated gonococcal infection

Total reports for BC is the sum of the following genders: female, male, transgender, and gender unknown

Perinatally-acquired Gonorrhoea

In 2014, there were no reports of perinatally-acquired gonorrhoea. From 2005 to 2014, two perinatal cases have been identified.

Gonorrhea Antimicrobial Resistance

Treatment of gonorrhea has long been challenged by the bacterium's ability to acquire resistance to multiple classes of antibiotics. Historically, effective antibiotics – penicillin, doxycycline, and ciprofloxacin – can no longer be used successfully against gonorrhea, leaving few remaining options. BC treatment guidelines currently recommend third-generation cephalosporins for the treatment of gonorrhea: injectable ceftriaxone (250mg) or oral cefixime (800mg), co-treated with 1g of azithromycin.¹³ Recent international surveillance data and case reports, however, suggest that susceptibility of gonorrhea to these current first-line treatments is also now threatened.^{14, 15} In this context, local surveillance is critical.

The BC Public Health Microbiology & Reference Laboratory (BCPHMRL) located at BCCDC routinely tests *N. gonorrhoeae* isolates for susceptibility to a panel of antimicrobial drugs, including cefixime, ceftriaxone, and azithromycin. Data presented here summarize the minimum inhibitory concentration (MIC) of these drugs among isolates from BC. The MIC is the lowest amount of antibiotic required to inhibit growth of the bacterium; a higher MIC means the bacterium is less susceptible to the antibiotic.




A total of 3,094 isolates were tested between 2006 and 2014, representing 23.0% (3,094/13,468) of all gonorrhea cases reported in the province. Fifty percent (1,525/3,094) of isolates tested for drug susceptibility were sampled from the urethra, 25.5% (788/3,094) from the rectum, 11.7% (362/3,094) from the cervix, and 12.3% (380/3,094) from the throat.

Since 2006, 0.6% (20/3,094) of isolates showed an MIC ≥ 0.25 $\mu\text{g}/\text{mL}$ to cefixime. Fortunately, no isolate was fully resistant to cefixime or ceftriaxone¹⁶ and no treatment failures were reported in BC during this period (Figure 21). The increasing trend in percentage of isolates with elevated MIC (i.e., reduced susceptibility) to cefixime or ceftriaxone observed in 2006-2010 reversed in 2011-2014. Similarly, the increasing trend in percentage of isolates with elevated MIC to azithromycin¹⁷ in 2006-2011 reversed in 2012-2014.

The decline observed in 2011-2014 for reduced susceptibility to cefixime or ceftriaxone among tested isolates is encouraging and, may in part be due to changes in the Canadian and provincial gonorrhea treatment guidelines to more effective regimens (i.e., increased cefixime dosage or improved medication adherence due to single dosage). These trends will be closely monitored in order to inform the future evolution of gonorrhea treatment recommendations. The continued threat of emerging resistance reinforces the need for STI prevention and control measures such as increased testing for gonorrhea, partner testing and treatment of gonorrhea, and tests of cure, as well as the need for antibiotic stewardship to ensure effective treatments for bacterial infections.

21. Percentage of tested *N. gonorrhoeae* isolates with elevated minimum inhibitory concentrations (MIC) to Cefixime, Ceftriaxone, and Azithromycin in BC, 2006 to 2014

Elevated MIC defined here as $\geq 0.064 \mu\text{g/mL}$ for cefixime/ceftriaxone and $\geq 0.5 \mu\text{g/mL}$ for azithromycin



Pelvic Inflammatory Disease and Ectopic Pregnancy

Pelvic inflammatory disease (PID) and ectopic pregnancy (EP) are medical conditions in women that can sometimes be caused by chlamydia or gonorrhea infection. Examination of the rates of these conditions can provide an indication of the burden of chlamydia and gonorrhea infections. Included in this report are data of physician billings (representing inpatient and outpatient treatment) and hospital discharges (representing inpatient treatment) provided by the BC Ministry of Health. Data are only presented up to 2013 due to lags in reporting, collation, and transfer of data.

In BC, steady declines in physician billings and hospital discharges related to PID have been observed over time. Rates in hospital discharges for EP have shown a similar trend although physician billings for EP have increased slightly in recent years. These trends, especially with the increase in chlamydia and gonorrhea infections among women, are encouraging that public health programs to improve detection and early treatment of these infections have been successful.¹⁸


Pelvic Inflammatory Disease

In 2013, the rate of physician billings related to PID decreased to 211.5 (1,924 physician billings) from 227.5 (2,069 physician billings) per 100,000 women aged 15-44 years in 2012 (Figure 22). Rate of hospital discharges related to PID show a decrease to 27.9 (254 hospital discharges) in 2013 from 31.8 (289 hospital discharges) per 100,000 women aged 15-44 years in 2012 (Figure 23).


Ectopic Pregnancy

The rate of physician billings related to EP increased to 219.1 (1,993 physician billings) in 2013 from 202.5 (1,841 physician billings) per 100,000 women aged 15-44 years in 2012 (Figure 22). In contrast, the rate of hospital discharges related to EP have decreased to 46.7 (425 hospital discharges) in 2013 from 49.7 (452 hospital discharges) per 100,000 women aged 15-44 years in 2012 (Figure 23).

22. Case reports of women aged 15-44 years with a physician billing related to pelvic inflammatory disease or ectopic pregnancy in BC, 1993 to 2013



23. Case reports of women aged 15-44 years with a hospital discharge related to pelvic inflammatory disease or ectopic pregnancy in BC, 1993 to 2013



Infectious Syphilis

Infectious Syphilis by Region, Gender, and Age


Syphilis infections are divided into several stages: primary, secondary, early latent, and late latent. The initial symptoms of syphilis may not always be recognized and without treatment individuals generally enter a prolonged asymptomatic phase. Individuals can still, however, be infectious despite not having any symptoms. Syphilis infection can lead to serious complications, including cardiovascular and neurologic disease, and may be fatal.

Following a decline in rates in BC in the early 1990's, infectious syphilis (i.e., primary, secondary, and early latent stages) began to re-emerge in BC starting in 1997, corresponding to a series of outbreaks in different populations. While provincial trends had been decreasing in 2009-2010, infectious syphilis rates began to increase in 2011 and this became more evident in 2012-2013. In BC, the rate of infectious syphilis decreased slightly in 2014 to 11.9 (549 cases) from 12.2 (558 cases) per 100,000 population in 2013 (Figure 24). The highest rate of infectious syphilis was in the Vancouver Coastal Health Authority (Figure 26). Across Health Service Delivery Areas, the highest rates were in Vancouver, Fraser North, and North Shore/Coast Garibaldi (Figure 25).


The majority of infectious syphilis cases in BC are male (Figure 27). Although male infectious syphilis rates decreased in 2009-2010, male infectious syphilis rates have since risen to 22.8 (524 cases) per 100,000 population in 2014. The highest rates among males were in those aged 25-39 years (Figure 31). In 2014, the rate of infectious syphilis among females was 1.0 (24 cases), a slight increase from 0.9 (20 cases) per 100,000 population in 2013 (Figure 27).

In 2014, 455 cases (82.9% of all cases) reported sexual (including anonymous) partners while the remaining cases (94 cases) provided no information regarding their sexual partners (83 cases) or were lost to follow-up (11 cases). Of these 455 cases, 357 cases (78.5%) reported 1,783 partners (the remaining 98 cases, 21.5% did not specify the number of partners). Almost 30% of cases (132/455) reported at least one anonymous partner.


24. Infectious syphilis case reports in BC and Canada, 1991 to 2014*




25. Infectious syphilis case reports in BC by health service delivery area, 2014



Southwestern BC Inset




Rate per 100,000 population by HSDA




ID	Health Service Delivery Area	Cases	Rate
11	East Kootenay	0	0.0
12	Kootenay Boundary	2	2.6
13	Okanagan	1	0.3
14	Thompson Cariboo Shuswap	4	1.8
21	Fraser East	7	2.4
22	Fraser North	55	8.6
23	Fraser South	34	4.4
31	Richmond	11	5.4
32	Vancouver	378	57.4
33	North Shore/Coast Garibaldi	17	6.0
41	South Vancouver Island	21	5.6
42	Central Vancouver Island	10	3.8
43	North Vancouver Island	1	0.8
51	Northwest	4	5.4
52	Northern Interior	1	0.7
53	Northeast	0	0.0

26. Infectious syphilis case reports in BC by health authority, 2005 to 2014




27. Infectious syphilis case reports in BC by gender, 2005 to 2014




* Other - transgender and gender unknown

28. Infectious syphilis case reports in BC by age group and gender, 2014




* Other - transgender and gender unknown


29. Infectious syphilis case reports in BC by age group - total, 2005 to 2014



30. Infectious syphilis case reports in BC by age group - female, 2005 to 2014



31. Infectious syphilis case reports in BC by age group - male, 2005 to 2014



Infectious Syphilis by Ethnicity

In males, the majority of cases in 2014 were among people who identified as Caucasian (282 cases; 53.8%) (Table 34). In comparison, the majority of cases in females in 2014 were among women who identified as Asian (9 cases, 37.5%), however, the trends are highly variable due to the small number of female cases each year (24 cases in 2014) (Table 33).

32. Percentage of infectious syphilis case reports in BC by ethnicity - total, 2005 to 2014

Ethnicity	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
<i>No. Diagnoses</i>	288	331	301	328	217	154	191	369	558	549
Caucasian	59.0	60.4	64.5	66.2	65.0	70.1	64.9	61.2	55.7	52.3
Aboriginal	15.6	12.4	10.6	8.8	7.8	3.9	3.7	5.7	3.9	2.9
Asian	12.2	11.5	8.6	10.7	6.5	12.3	9.9	10.6	12.2	13.5
South Asian	3.8	6.0	4.0	4.3	1.4	3.2	6.8	4.1	3.6	2.6
Hispanic	5.6	4.2	5.6	4.9	9.2	2.6	8.4	6.2	5.2	6.0
Black	2.1	2.7	2.3	2.4	4.6	1.3	0.5	1.1	1.6	2.4
Other*	0.3	2.4	2.0	1.2	3.7	2.6	3.1	1.9	1.3	1.5
Unknown	1.4	0.3	2.3	1.5	1.8	3.9	2.6	9.2	16.5	18.9

* Other - Arab/West Asian, South American, and other/mixed ethnicity

33. Percentage of infectious syphilis case reports in BC by ethnicity - female, 2005 to 2014

Ethnicity	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
<i>No. Diagnoses</i>	86	77	55	41	28	13	13	24	20	24
Caucasian	39.5	40.3	54.5	43.9	57.1	69.2	38.5	29.2	15.0	20.8
Aboriginal	37.2	42.9	34.5	29.3	28.6	15.4	15.4	20.8	15.0	8.3
Asian	11.6	7.8	5.5	12.2	0.0	0.0	23.1	16.7	30.0	37.5
South Asian	1.2	6.5	3.6	9.8	7.1	7.7	7.7	12.5	10.0	4.2
Hispanic	5.8	0.0	0.0	0.0	0.0	0.0	7.7	0.0	5.0	0.0
Black	1.2	1.3	0.0	2.4	3.6	7.7	0.0	4.2	0.0	0.0
Other*	0.0	1.3	0.0	0.0	3.6	0.0	0.0	0.0	0.0	0.0
Unknown	3.5	0.0	1.8	2.4	0.0	0.0	7.7	16.7	25.0	29.2

* Other - Arab/West Asian, South American, and other/mixed ethnicity

34. Percentage of infectious syphilis case reports in BC by ethnicity - male, 2005 to 2014

Ethnicity	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
<i>No. Diagnoses</i>	202	254	245	286	188	141	178	344	536	524
Caucasian	67.3	66.5	66.9	69.6	66.5	70.2	66.9	63.4	57.3	53.8
Aboriginal	6.4	3.1	5.3	5.6	4.8	2.8	2.8	4.7	3.5	2.7
Asian	12.4	12.6	9.4	10.5	7.4	13.5	9.0	10.2	11.6	12.4
South Asian	5.0	5.9	4.1	3.5	0.5	2.8	6.7	3.5	3.4	2.5
Hispanic	5.4	5.5	6.5	5.6	10.6	2.8	8.4	6.7	5.2	6.3
Black	2.5	3.1	2.9	2.4	4.8	0.7	0.6	0.9	1.7	2.5
Other*	0.5	2.8	2.4	1.4	3.2	2.8	3.4	2.0	1.1	1.5
Unknown	0.5	0.4	2.4	1.4	2.1	4.3	2.2	8.7	16.2	18.3

* Other - Arab/West Asian, South American, and other/mixed ethnicity


Infectious Syphilis among Aboriginal Peoples

This section describes cases of infectious syphilis among people who identify as Aboriginal. Among the nearly 200,000 Aboriginal persons living in BC, approximately 66% are First Nations, 30% are Métis, and fewer than 5% are Inuit or of other Aboriginal identity.¹⁹

Since 2005, the percentage of infectious syphilis cases in BC residents that were reported among Aboriginal peoples has been decreasing. In 2014, the percentage of infectious syphilis cases among Aboriginal peoples decreased to 2.9% (16 cases) from 3.9% (22 cases) in 2013 (Table 32). In comparison, Aboriginal peoples represent about 5% of the general BC population.²⁰ While numbers are small, Aboriginal women in BC are disproportionately represented among all female infectious syphilis cases (2 cases or 8.3% of all female cases in 2014).

Consistent with overall provincial trends, the rate of infectious syphilis among First Nations people in BC has increased since 2010 (Figure 35). The rate decreased in 2014 to 11.4 (16 cases) from 15.1 (21 cases) per 100,000 population in 2013. Due to limitations in the availability of population estimates of Métis and Inuit peoples, it is not possible to calculate comparable rates. (See Technical Appendix for further details about the classification of ethnicity for syphilis cases and First Nations population estimates.)

35. Infectious syphilis case reports among First Nations people in BC by gender, 2005 to 2014



Rates based on First Nations population estimates from Aboriginal Affairs and Northern Development Canada (AANDC)

Infectious Syphilis by Exposure Category

Gay, bisexual, and other men who have sex with men (MSM) continue to comprise the greatest number of infectious syphilis cases in BC (Figure 37). The number of syphilis cases among MSM decreased slightly from 476 cases (85.3% of all cases) in 2013 to 466 cases (84.9%) in 2014. Infectious syphilis cases among heterosexual persons without other risk factors plateaued in 2013-2014 (51 cases, 9.1% in 2013; 51 cases, 9.3% in 2014). The number of infectious syphilis cases among street involved persons, sex trade workers, and their patrons has continued to decrease (14 cases, 2.5% in 2013; 6 cases, 1.1% in 2014). Trends among MSM in BC are explored in more detail in the next section.

36. Infectious syphilis case reports in BC by exposure category and health authority, 2005 to 2014

Health Authority	Exposure Category	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Interior	MSM	1	2	7	2	5	1	3	6	4	3
	Street/STW	0	1	3	3	1	1	0	0	0	0
	HET	3	2	0	2	7	2	4	2	2	2
	Outside	0	0	0	0	0	0	0	1	0	1
	Other/UNK	0	0	0	0	0	0	0	0	0	1
Fraser	MSM	10	22	13	41	25	17	15	46	70	78
	Street/STW	23	35	23	19	4	2	2	3	4	1
	HET	22	25	21	17	13	5	7	11	18	11
	Outside	3	3	3	4	1	2	0	1	2	2
	Other/UNK	0	1	1	1	0	0	3	3	4	4
Vancouver Coastal	MSM	108	129	136	179	108	94	121	248	386	356
	Street/STW	71	62	34	23	8	7	6	8	5	5
	HET	37	27	15	14	18	13	6	17	29	29
	Outside	2	6	5	1	2	3	2	0	2	4
	Other/UNK	1	0	1	1	1	1	2	6	9	12
Vancouver Island	MSM	1	4	7	6	6	3	13	10	13	22
	Street/STW	1	0	3	3	4	0	0	0	0	0
	HET	2	8	7	2	8	1	3	2	6	8
	Outside	0	1	0	0	0	0	0	0	0	1
	Other/UNK	0	0	0	1	1	0	0	1	1	1
Northern	MSM	0	0	6	1	0	0	1	0	1	4
	Street/STW	0	2	2	4	1	0	1	1	0	0
	HET	1	1	8	4	2	2	2	3	0	1
	Outside	0	0	0	0	0	0	0	0	0	0
	Other/UNK	0	0	0	0	0	0	0	0	1	0

MSM - men who have sex with men


Street/STW - street involved / sex trade worker or patron

HET - heterosexual contact


Outside - acquired outside of Canada

UNK - exposure unknown


37. Infectious syphilis case reports in BC by exposure category - total, 2005 to 2014



38. Infectious syphilis case reports in BC by exposure category - female, 2005 to 2014



39. Infectious syphilis case reports in BC by exposure category - male, 2005 to 2014



Infectious Syphilis among Men who have Sex with Men

Gay, bisexual, and other men who have sex with men (MSM) continue to be disproportionately affected by the current infectious syphilis epidemic in BC, constituting 84.9% (466/549 cases) of all cases in 2014 (Figure 37). The number of infectious syphilis cases among MSM has been steadily increasing since 2011. This is similar to the profile of syphilis epidemics in other Canadian provinces²¹, the US²², and several European countries²³ where syphilis cases were also predominantly among MSM and trends are increasing.

There were 466 cases among MSM in 2014, 15.5% (72 cases) were diagnosed with primary syphilis, 19.1% (89 cases) with secondary syphilis, and 65.5% (305 cases) were diagnosed with early latent infection. The proportion of infectious syphilis cases among MSM in the early latent stage of infection has steadily increased over time which may reflect increased syphilis testing. In 2014, 112 (24.0%) MSM cases had a prior syphilis diagnosis within the past five years, highlighting the importance of repeat infections in the current epidemic.²⁴

Despite recent increases in the number of annual cases, the characteristics of MSM with syphilis have remained similar over time. While the number of cases has decreased for most age groups in 2014 (Figure 41), the mean age of MSM diagnosed with syphilis has remained relatively stable at 42 years (range 19-84 years). In 2014, the majority of cases resided in the Lower Mainland with 76.4% (356 cases) residing in Vancouver and 16.7% (78 cases) in Fraser. As in previous years, the majority of cases in 2014 were among Caucasian (258 cases, 55.4%), Asian (61 cases, 13.1%), and Hispanic (34 cases, 7.3%) men (Table 40).


Being HIV positive continues to be an important risk factor for acquiring infectious syphilis. In 2014, of the MSM cases with a known HIV status (447 cases), 59.7% (267 cases) were HIV positive at the time of their syphilis diagnosis which is a decrease from 2013 (305/473 cases, 64.5%) (Figure 42). The possible roles of core sexual networks and the biological synergy between HIV and syphilis are important areas of study which may help explain this trend. Public health follow-up, partner notification, and partner testing for all syphilis cases remain the cornerstone to controlling the syphilis epidemic for MSM in BC as do efforts to raise awareness among MSM. Given the continued increase of syphilis cases, the enhancement of ongoing programs and development of new syphilis control interventions for MSM remain a priority in BC.

40. Percentage of infectious syphilis case reports among MSM in BC by ethnicity, 2005 to 2014


Ethnicity	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
<i>No. Diagnoses</i>	122	157	174	229	146	115	154	310	476	466
Caucasian	77.9	79.6	73.0	75.5	68.5	75.7	70.1	64.8	59.9	55.4
Aboriginal	3.3	0.6	2.3	2.6	3.4	3.5	1.9	4.5	3.6	2.6
Asian	10.7	7.6	8.6	10.0	5.5	7.8	7.1	9.4	10.5	13.1
South Asian	3.3	3.2	1.7	3.1	0.0	3.5	5.8	3.2	2.7	2.4
Hispanic	1.6	6.4	8.0	4.8	13.0	3.5	9.7	7.1	5.7	7.3
Black	1.6	1.3	3.4	1.3	3.4	0.0	0.0	1.0	1.7	2.1
Other*	0.8	1.3	2.3	1.7	4.1	2.6	3.2	2.3	1.5	1.1
Unknown	0.8	0.0	0.6	0.9	2.1	3.5	1.9	7.7	14.5	16.1

* Other - Arab/West Asian and other/mixed ethnicity

41. Infectious syphilis case reports among MSM in BC by age group, 2005 to 2014



42. Infectious syphilis case reports* among MSM in BC by HIV co-infection, 2005 to 2014




Stage of Infection at Time of Syphilis Diagnosis

There have been slow changes in the proportion of infectious syphilis cases by stage of infection (i.e., primary, secondary, and early latent) over time. Most notably, the proportion of cases that are early latent syphilis has been increasing over time. This may reflect a greater uptake of syphilis testing or screening as people with syphilis infections in the early latent stage of infection are typically asymptomatic. In 2014, the proportion of cases that are early latent increased to 65.4% (359 cases) from 55.6% (310 cases) in 2013 (Figure 43). This increase may be a year-to-year variation.

From 2005 to 2014, 5.3% (175/3,286 cases) of infectious syphilis cases were also diagnosed with neurosyphilis. Neurosyphilis is commonly considered to be indicative of an advanced syphilis infection but can sometimes occur at an early stage of infection.

43. Stage of infection at time of syphilis diagnosis, 2005 to 2014




* Neurosyphilis is not mutually exclusive from other stages of syphilis infection (i.e., primary, secondary and early latent)

Maternal and Early Congenital Syphilis

Infectious syphilis acquired prior to or during pregnancy can be passed from a mother to her infant which can have serious consequences, including infant death. Accordingly, prenatal screening for syphilis is recommended for all pregnant women in BC, as treatment will reduce the risk of transmission to or complications in infants.

In 2014, 50,741 syphilis tests were conducted as part of prenatal screening. Overall, the number of syphilis tests performed prenatally each year in BC is increasing slightly. There were no cases of congenital syphilis identified in BC in 2014 (Figure 44). In 2014, six maternal syphilis cases were reported whereas seven cases were reported in 2013.

44. Maternal and early congenital syphilis case reports in BC, 2005 to 2014



Reports - Maternal	16	14	8	14	13	5	3	8	7	6
Reports - Early Congenital	3	2	2	2	2	0	0	0	1	0
Rate * - Early Congenital	7.4	4.8	4.6	4.5	4.5	0.0	0.0	0.0	2.3	0.0
Total number of syphilis prenatal tests			45,016	47,098	47,011	47,348	48,726	50,040	49,724	50,741

* Rate per 100,000 live births

Endnotes

- 1 For more information about the increase in chlamydia infections in high income countries around the world see: European Centre for Disease Prevention and Control. (February 2014). Chlamydia control in Europe: literature review. Retrieved from <http://www.ecdc.europa.eu/en/publications/Publications/chlamydia-control-europe.pdf>
- 2 Reference to the increase in chlamydia infections in high income countries around the world see: Centers for Disease Control and Prevention, Division of STD Prevention. (January 2014). Sexually Transmitted Disease Surveillance 2012. Retrieved from <http://www.cdc.gov/std/stats12/Surv2012.pdf>
- 3 Reference to the “arrested immunity” hypothesis: Brunham RC, Rekart ML. (January 2008). The arrested immunity hypothesis and the epidemiology of chlamydia control. *Sexually Transmitted Diseases*, 35(1), 53-54.
- 4 Reference to the “arrested immunity” hypothesis: Rekart ML, Gilbert M, Meza R, Kim PH, Chang M, Money DM et al. (2012). Chlamydia public health programs and the epidemiology of pelvic inflammatory disease and ectopic pregnancy. *Journal of Infectious Diseases*, advance access published October 24, 2012. Retrieved from <http://jid.oxfordjournals.org/content/early/2012/10/24/infdis.jis644.full.pdf?keytype=ref&ijkey=y7YveY9ycdjbCr2>
- 5 For information about the increase in lymphogranuloma venereum (LGV) in the US: Schachter J, Moncada J. (June 2005). Lymphogranuloma venereum: how to turn an endemic disease into an outbreak of a new disease? Start looking. *Sexually Transmitted Diseases*, 32(6), 331-332. Retrieved from http://journals.lww.com/stdjournal/Citation/2005/06000/Lymphogranuloma_Venereum__How_to_Turn_an_Endemic.1.aspx
- 6 Reference to increase in lymphogranuloma venereum (LGV) in Europe: Nieuwenhuis RF, Ossewaarde JM, Götz HM, Dees J, Thio HB, Thomeer MGJ et al. (2004). Resurgence of Lymphogranuloma venereum in Western Europe: an outbreak of chlamydia trachomatis serovar L2 proctitis in the Netherlands among men who have sex with men. *Clinical Infectious Diseases*, 39(7), 996-1003. Retrieved from <http://cid.oxfordjournals.org/content/39/7/996.full.pdf+html>
- 7 For more information about lymphogranuloma venereum (LGV) in BC: BC Centre for Disease Control. (March 2012). Lymphogranuloma venereum in British Columbia, 2004 to 2011. Retrieved from http://www.bccdc.ca/NR/rdonlyres/27E4F543-7D0E-417B-AF41-AEB262776FC3/0/STI_Reports_LGVinBC_20120404.pdf
- 8 For more information about lymphogranuloma venereum (LGV) in BC: BC Centre for Disease Control. (June 12, 2015). Lymphogranuloma venereum in BC: An update on a re-emerging STI. Retrieved from http://www.bccdc.ca/NR/rdonlyres/68FC8661-6622-42BE-8C71-F315382866E9/0/CPS_LGV_Update_20150612.pdf
- 9 For information about the how an individual with gonorrhea is at increased risk of acquiring HIV see: Wilton J. (Spring 2012). STIs: What role do they play in HIV transmission? CATIE, Canada’s source for HIV and hepatitis C information. Retrieved from <http://www.catie.ca/en/pif/spring-2012/stis-what-role-do-they-play-hiv-transmission>
- 10 Reference to how an individual with gonorrhea is at increased risk of acquiring HIV: Bernstein KT, Marcus JL, Nieri G, Philip SS, Klausner JD. (April 1, 2010). Rectal gonorrhea and chlamydia reinfection is associated with increased risk of HIV seroconversion. *Journal of Acquired Immune Deficiency Syndromes*, 53(4), 537-543. Retrieved from <http://journals.lww.com/jaids/toc/2010/04010>
- 11 Reference to the increase in gonorrhea infections among MSM in other jurisdictions: Public Health England. (June 20, 2014). Sexually transmitted infections and chlamydia screening in England, 2013. Health Protection Report, 8(24). Retrieved from https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/326935/hpr2414.pdf
- 12 Reference to the increase in gonorrhea infections among MSM in other jurisdictions: Wolitski RJ, Fenton KA. (2011). Sexual health, HIV, and sexually transmitted infections among gay, bisexual, and other men who have sex with men in the United States. *AIDS and Behavior*, 15 Suppl 1:S9-17.
- 13 For more information about the BC treatment guidelines for gonorrhea see: BC Centre for Disease Control. (revised August 2014). British Columbia Treatment Guidelines: Sexually Transmitted Infections in Adolescents and Adults 2014. Retrieved from http://www.bccdc.ca/NR/rdonlyres/46AC4AC5-96CA-4063-A563-0BA9F4A0A6E9/0/CPS_BC_STI_Treatment_Guidelines_27082014.pdf
- 14 Reference to international surveillance data that suggests susceptibility of gonorrhea to cephalosporin treatments is threatened: Unemo M, Golparian D, Syversen G, Vestrheim DF, Moi H. (November 25, 2010). Two cases of verified clinical failures using internationally recommended first-line cefixime for gonorrhoea treatment, Norway, 2010. *Euro Surveillance*, 15(47). Retrieved from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19721>

- ¹⁵ Reference to international surveillance data that suggests susceptibility of gonorrhoea to cephalosporin treatments is threatened: Yokoi S, Deguchi T, Ozawa T, Yasuda M, Ito S, Kubota Y, et al. (August 2007). Threat to cefixime treatment for gonorrhoea. *Emerging Infectious Diseases*, 13(8), 1275-1277. Retrieved from: <http://wwwnc.cdc.gov/eid/content/13/8/pdfs/v13-n8.pdf>
- ¹⁶ Minimum inhibitory concentration (MIC) breakpoints to define “resistance” to cefixime and ceftriaxone have not yet been established, however, the Clinical and Laboratory Standards Institute (CLSI) defines MIC \leq 0.25 $\mu\text{g}/\text{mL}$ as susceptible.
- ¹⁷ The US Centers for Disease Control and Prevention (CDC) has proposed MIC \geq 2 $\mu\text{g}/\text{mL}$ as the non-susceptible threshold for azithromycin.
- ¹⁸ See Endnote #3.
- ¹⁹ BC Stats. Census Statistical Profiles of Aboriginal Peoples, 2006. Retrieved from <http://www.bcstats.gov.bc.ca/statisticsbysubject/AboriginalPeoples/CensusProfiles.aspx>
- ²⁰ See Endnote #19
- ²¹ Reference to the increase in infectious syphilis cases among MSM in Canada: Public Health Agency of Canada. (2009). Executive Summary - Report on sexually transmitted infections in Canada: 2009. Retrieved from <http://www.phac-aspc.gc.ca/sti-its-surv-epi/sum-som-eng.php>
- ²² Reference to the increase in infectious syphilis cases among MSM in the US: US Center for Disease Control and Prevention. (2011). 2011 Sexually Transmitted Diseases Surveillance – Syphilis. Retrieved from <http://www.cdc.gov/std/stats11/syphilis.htm>
- ²³ Reference to the increase in infectious syphilis cases among MSM European countries: Savage EJ, Hughes G, Ison C, Lowndes CM, the European Surveillance of Sexually Transmitted Infections (ESSTI) network. (November 26, 2009). Syphilis and gonorrhoea in men who have sex with men: a European overview. *Euro Surveillance*, 14(47). Retrieved from <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19417>
- ²⁴ For more information about the increase in infectious syphilis among MSM in BC: BC Centre for Disease Control. (June 2013). Infectious syphilis among gay, bisexual and other men who have sex with men in British Columbia, 2003 to 2012. Retrieved from http://www.bccdc.ca/NR/rdonlyres/B917A2F2-54C5-4691-8015-4F80538CAC1E/0/CPS_Report_Infectious_Syphilis_MSMBC_20032012_20130624.pdf
- ²⁵ Clinical presentation includes any evidence of congenital syphilis on physical examination (e.g., hepatosplenomegaly), evidence of congenital syphilis on radiographs of long bones, a reactive CSF VDRL, an elevated CSF cell count or protein without other cause. Note that neonates may not display clinical manifestations of congenital syphilis and may meet laboratory criteria only.

Contributors

Epidemiology & Surveillance Team Clinical Prevention Services

Dr. Jason Wong, Physician Epidemiologist
Elsie Wong, Federal Field Surveillance Officer (PHAC)
Sung Jae Lee, Surveillance Analyst
Dr. Theodora Consolacion, Epidemiologist

We would like to acknowledge the contributions of our many partners who without their support this report would not have been possible.

- Staff from the BC Public Health Microbiology & Reference Laboratory (BCPHMRL), located at BCCDC, for the collecting and compiling of gonorrhea antibiotic susceptibility data.
- Physicians, health care providers, and public health staff in BC for taking the time and effort to complete and submit case report forms.
- Staff from Clinical Prevention Services, BCCDC for the collecting (Epid and Syphilis Nurses) and entering (Clerical Team) of STI data.
- First Nations Health Authority for providing feedback to the section pertaining to First Nations people.
- BC Ministry of Health for providing pelvic inflammatory disease and ectopic pregnancy data.
- Surveillance and Epidemiology Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada for providing the national chlamydia, gonorrhea and infectious syphilis rates.

Technical Appendix

Data Limitations

There are several key limitations to surveillance data which are important to understand in order to interpret surveillance data appropriately.

- Surveillance data are only reflective of the proportion of the population who test. Individuals with infections who have not tested are not included in the surveillance data presented in this report. Many people with sexually transmitted infections do not have symptoms and so do not seek health care advice or testing. This under-counting of cases may disproportionately affect some groups. For example, more women than men get syphilis testing in BC because of prenatal syphilis screening. Also, some sexually transmitted infections are more or less likely to show symptoms depending on sex and site of infection (e.g., urethral gonococcal infections in men are more likely to produce symptoms than those in women or those in the pharynx/throat).
- Surveillance trends are influenced by provider testing behaviours which may result in changes to the number of tests performed each year (e.g., increased vigilance for lymphogranuloma venereum in 2010-2011 may have resulted in more diagnoses of this disease, artificially driving up lymphogranuloma rates).
- Trends are influenced by temporal changes in testing technologies. Over the past ten years, nucleic acid amplification tests (NAAT) have replaced culture-based diagnostics for chlamydia and gonorrhoea testing. The use of NAAT, which is a more sensitive test, has resulted in increased detection of these infections. NAAT technology has also allowed urine-based screening for chlamydia and gonorrhoea thus reducing the use of urethral swabs for asymptomatic male clients. With the use of this less-invasive procedure, more men may be getting screened for chlamydia and gonorrhoea.
- The majority of surveillance data presented in this surveillance report are extracted from case report forms completed by health care providers or public health nurses as part of the case follow-up process (which includes partner notification, patient education, and referral to appropriate services). There is an expected reporting delay to receipt of these forms which may range from days to months depending on the infection.
- Cases are classified by exposure category and ethnicity according to information elicited during follow-up from the case or their health care provider and, under-reporting of this information may lead to misclassification. For example, gay, bisexual and other men who have sex with men, and transgender persons may be reluctant to disclose these factors to their health care providers due to social stigma.
- Rates of infectious syphilis diagnoses among First Nations people are calculated with the numerator comprised of individuals with infectious syphilis who self identify as First Nations and the denominator comprised of individuals who are registered First Nations (see Data Sources below for further details).

Case Definitions

Sexually transmitted infections (STIs) included in this report are listed as reportable diseases in the *Communicable Disease Regulation* (Schedule A) of the *Public Health Act*.

Chlamydia

Genital: Detection and confirmation of *C. trachomatis* in anogenital (including rectal) or urinary specimens by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen).

Extra-genital: Detection and confirmation of *C. trachomatis* in specimens from the conjunctiva, pharynx and other extra-genital sites by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen).

Perinatally-acquired: Detection and confirmation of *C. trachomatis* in nasopharyngeal or other respiratory tract specimens by appropriate laboratory techniques (e.g., isolation of *C. trachomatis* by culture, demonstration of *C. trachomatis* nucleic acid or antigen) from an infant who developed pneumonia in the first 6 months of life or conjunctival specimens from an infant who developed conjunctivitis in the first month of life.

Gonorrhoea

Genital: Detection and confirmation of *N. gonorrhoeae* in anogenital (including rectal) or urinary specimens by appropriate laboratory techniques (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Extra-genital: Detection and confirmation of *N. gonorrhoeae* in specimens from the conjunctiva, pharynx, joint, blood, and other extra-genital sites by appropriate laboratory techniques (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Perinatally-acquired: Detection and confirmation of *N. gonorrhoeae* infection in the first 4 weeks of life leading to the diagnosis of gonococcal conjunctivitis, scalp abscess, vaginitis, bacteremia, arthritis,

meningitis, or endocarditis by an appropriate laboratory technique (e.g., culture, detection of *N. gonorrhoeae* nucleic acid).

Infectious Syphilis

Infectious syphilis is a complex sexually transmitted infection that has a highly variable clinical course. Three stages of syphilis, primary, secondary and early latent, comprise infectious syphilis. Classification by a clinician with expertise in syphilis may take precedence over the following case definitions developed for surveillance purposes.

Primary Syphilis: Current clinical presentation compatible with primary syphilis (e.g., one or more ulcers/chancres), and one of the following:

- Identification of *T. pallidum* in clinical specimens (e.g., from chancre, regional lymph node) by dark field microscopy, direct fluorescent antibody, or nucleic acid amplification test (NAAT), or
- Reactive serology (treponemal, regardless of non-treponemal serology reactivity) in individuals with no previous history of syphilis, or
- Significant (i.e., four-fold or greater) increase in titre over the last known non-treponemal test.

Secondary Syphilis: Clinical presentation compatible with secondary syphilis (e.g., rash, fever, malaise, lymphadenopathy, mucus lesions, condyloma lata, alopecia, meningitis, headaches, uveitis, retinitis, recent hearing impairment), and one of the following:

- Identification of *T. pallidum* in clinical specimens (e.g., from chancre, regional lymph node) by dark-field microscopy, direct fluorescent antibody, or nucleic acid amplification test (NAAT), or

- Reactive serology (non-treponemal and treponemal) in individuals with no previous history of syphilis, or
- Significant (i.e., four-fold or greater) increase in titre over the last known non-treponemal test.

Early Latent Syphilis: An individual without symptoms of primary or secondary syphilis but has a reactive serology (non-treponemal and treponemal) or four-fold increase in titre over the last known non-treponemal test, and one of the following within the previous 12 months:

- Non-reactive serology, or
- Symptoms suggestive of primary or secondary syphilis, or
- Exposure to a sexual partner with primary, secondary or early latent syphilis, or
- Is a member of (or has had sexual partners in the previous 12 months from) groups at known increased risk of syphilis infection in BC, or
- Has a titre of $\geq 1:16$.

Early Congenital Syphilis: A stillbirth, neonate or older individual with clinical presentation²⁵ compatible with congenital syphilis, onset less than two years of age, and one of the following:

- Four-fold higher RPR than maternal titre and positive treponemal confirmatory test, or
- Detection of *T. pallidum* in clinical specimens (e.g., lesions, placenta, umbilical cord, autopsy) through darkfield microscopy, direct fluorescent antibody assay, or PCR, or
- Mother with untreated or inadequately treated syphilis (primary, secondary, early or late latent syphilis) during pregnancy or at birth.

Maternal Syphilis: A woman who meets the case definition of infectious syphilis (primary, secondary, early latent syphilis) or late latent syphilis, and one of the following:

- Syphilis serology conducted as part of prenatal blood screening, or
- Known to have given birth to an infant (live or stillborn) with congenital syphilis, or

- Clinical presentation with infectious syphilis during pregnancy.

Lymphogranuloma Venereum (LGV)

Confirmed: DNA sequencing for *C. trachomatis* confirming serovars of L1, L2, or L3 present.

Probable: One of the following two case definitions:

- Positive chlamydia NAAT or culture (from any site), and one of the following:
 - Proctitis, or
 - Inguinal/femoral lymphadenopathy, or
 - Suspicious lesion, or
 - Sexual partner who is confirmed or probable LGV case
- Clinical symptoms consistent with LGV (proctitis or inguinal/femoral lymphadenopathy or suspicious lesion) without a positive chlamydia test, and sexual partner who is confirmed or probable LGV case.

Data Sources

STI Data (Chlamydia, Gonorrhea, Infectious Syphilis)

When an individual is diagnosed with a reportable STI, the care provider completes a case report form (Health 208 form) then forwards it to BCCDC where the information is entered into the provincial STI database. Public health clinics with access to the provincial STI database directly enter the information for their newly diagnosed individuals.

Pelvic Inflammatory Disease and Ectopic Pregnancy Data

The diagnoses of pelvic inflammatory disease (PID) and ectopic pregnancy (EP) are captured in the Discharge Abstract Database (DAD) and the Medical Service Plan (MSP) payment database maintained by the BC Ministry of Health. The DAD includes data on patient discharges and day surgeries directly from hospitals in BC, including all known facilities for acute care and day surgery, and most facilities for

chronic care and rehabilitation. The MSP database contains data on insured medical services delivered on a fee-for-service basis. This data includes physician billings for inpatient and outpatient care, claims from supplementary health care practitioners, and claims for laboratory services and diagnostic procedures. MSP data do not include services provided via an Alternate Payment Program (e.g., contract or salary). Approximately 30% of BC physicians receive some remuneration through alternative payments. In particular, physicians in rural areas are more likely to be paid with Alternate Payment Programs. Some hospital emergency departments in BC are also funded through Alternate Payment Program contracts. Administrative data on PID and EP included in this surveillance report are extracted for women of reproductive age (15-44 years) who have at least one physician billing or hospital discharge per year based on the following International Classification of Disease (ICD) codes:

Pelvic Inflammatory Disease

ICD 9: MSP (1992-2010), DAD (1992-2000)

- Salpingitis and oophoritis (614, 614.0-614.2)
- Parametritis and pelvic cellulitis/peritonitis (614.3-614.5, 614.7)
- Other or unspecified inflammatory disease of female pelvic organs and tissues (614.8, 614.9)
- Inflammatory diseases of uterus except cervix (615, 615.0-615.9)

ICD 10: DAD (2001-2010)

- Salpingitis and oophoritis (N70, N70.0-N70.9)
- Parametritis and pelvic cellulitis/peritonitis (N73.0-N73.5)
- Other or unspecified female pelvic inflammatory diseases (N73.8, N73.9)
- Inflammatory diseases of uterus except cervix (N71, N71.0-N71.9)

Ectopic Pregnancy

ICD 9: MSP (1992-2010), DAD (1992-2000)

- Ectopic pregnancy (633)
- Abdominal pregnancy (633.0)

- Tubal pregnancy (633.1)
- Ovarian pregnancy (633.2)
- Other or unspecified ectopic pregnancy (633.8, 633.9)

ICD 10: DAD (2001-2010)

- Ectopic pregnancy (000)
- Abdominal pregnancy (000.0)
- Tubal pregnancy (000.1)
- Ovarian pregnancy (000.2)
- Other or unspecified ectopic pregnancy (000.8, 000.9)

BC Public Health Microbiology & Reference Laboratory located at BCCDC

The BC Public Health Microbiology & Reference Laboratory (BCPHMRL) performs approximately 15-20% of all gonorrhea testing in the province, receiving specimens predominantly from Provincial Sexually Transmitted Infection Clinic sites operated by the BCCDC, from regional public health, youth, reproductive and sexual health clinics, and from hospitals throughout the province. At BCPHMRL, gonorrhea may be detected by nucleic acid amplification testing (NAAT) or conventional culture diagnostic methods. Culture testing is preferentially used for rectal and pharyngeal specimens and for all specimens from contacts to gonorrhea as well as patients who are symptomatic, not responding to treatment, or presenting for treatment after an initial NAAT-positive test. Antimicrobial susceptibility testing is routinely performed for all *N. gonorrhoeae* isolated by culture from clinical specimens. The BCPHMRL additionally receives gonorrhea isolates forwarded for susceptibility testing from community or hospital-based laboratories in BC. Antimicrobial susceptibility testing is by E-test (bioMerieux) and data are analyzed by isolate.

Population Data

Unless noted otherwise, population data and associated rates are based on the P.E.O.P.L.E. 2014 Population Estimates and Projections released by BC Stats, BC Ministry of Labour and Citizens' Services.

First Nations Population Estimates

Population rates for First Nations people are calculated using estimates from Aboriginal Affairs and Northern Development Canada.

These estimates are based on the Indian Registry System (IRS), which includes individuals who have registered for First Nations status under the Indian Act. The IRS is subject to several limitations, including:

- Under-counting due to delayed reporting of infants entitled to be registered, as well as other unregistered individuals who are entitled for status designation
- Over-counting due to individuals remaining on the IRS after they are deceased
- Geographic misclassification because individuals are included in the BC population according to membership of a BC band rather than current place of residence
- Systematic biases from imbalance in the migration into and out of the BC region (these are difficult to quantify)

For further details about the data source and its limitations, see the report entitled *Registered Indian Population by Sex and Residence, 2014: Statistics and Measurement Directorate (2014)*. Aboriginal Affairs and Northern Development Canada.

Live Births

From 2005-2012, perinatal rates are calculated using live births data from the BC Vital Statistics Agency and, for 2013 live births data from Perinatal Service BC – British Columbia Perinatal Data Registry (data provided on September 11, 2015).

Additional Notes

Classification of Health Region

Cases are assigned to health regions (i.e., Health Authority or Health Service Delivery Area) by residence. If residence is unknown, the case is then assigned to the health region where the individual was tested.

Classification of Ethnicity

Ethnicity is based on information elicited from the case or health care provider during follow-up. Since ethnicity data for chlamydia and gonorrhoea cases are often not collected they are not included in this report.

Ethnicity	Example
Aboriginal*	First Nations, Inuit, Métis
Arab/West Asian	Aremnian, Egyptian, Iranian, Moroccan, Lebanese, Afghani
Asian	Chinese, Japanese, Vietnamese, Cambodian, Indonesian, Filipino, Korean, Laotian
Black	African, Haitian, Jamaican, Somali
Caucasian (White)	Irish, Scottish, English, Portuguese, Italian, Russian
Hispanic	Mexican, Central/South American
South Asian	East Indian, Pakistani, Sri Lankan, Punjabi, Bangladeshi
other/mixed ethnicity	ethnicity is known but is not included in one of the above categories or case has dual ethnicity
unspecified	information about ethnicity is not elicited from case or health care provider

* For infectious syphilis, enhanced case report forms have not included response categories for Inuit or Métis. On this basis, some Métis individuals may have been incorrectly described as First Nations in these surveillance data. Revisions to reporting forms are underway to correct this issue.

Exposure Group Hierarchy

Cases may have more than one type of sexual exposure. The following are definitions of sexual exposures used in this surveillance report. For infectious syphilis cases, individuals are assigned to the exposure category listed first (or highest) in the following hierarchy.

1. **MSM***: Male who reports having male sex partner(s), with or without female sex partners.

2. Street-Involved, Sex Trade Worker and Patron:

- i) Street-Involved – Person who reports either: (a) living on the street or in a single room occupancy (SRO) hotel; or (b) attached to the street; or (c) having no fixed address; or (d) transient
- ii) Sex Trade Worker (STW) – Person who reports providing sex to another individual in exchange for money, shelter, food, drugs, etc.
- iii) Patron of STW – Person who reports payment (with money, shelter, food, drugs, etc.) for sex with a STW

3. Heterosexual Contact*: Male who reports having female sex partner(s) only or female who reports having male with/without female sex partner(s).

4. Acquired Outside of Canada:

- i) Foreign Acquired – Person currently residing in Canada but likely acquired syphilis outside of Canada (i.e., reports sexual partner(s) in other countries)
- ii) Immigration – Person immigrating to Canada and identified with syphilis through testing done as part of the immigration process

5. Other Risk Factor: Likely route of exposure is known but cannot be classified into any of the major exposure categories listed here. For example, females reporting female sex partner(s) only.

6. Unknown: Route of exposure is unknown or not identified at the time of completion of case follow-up (e.g., route of exposure not provided by case).

* A transgender individual may be assigned to either MSM or Heterosexual Contact exposure category depending on how this individual describes their sexual partners.