

BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN

2008-09: Number 33, Week 37

September 13 – 19, 2009



BC Centre for Disease Control
An agency of the Provincial Health Services Authority

Prepared by BCCDC Influenza &
Emerging Respiratory Pathogens Team

Increasing Influenza Activity in BC, Predominantly Attributed to Pandemic pH1N1

Contents:

British Columbia:

Sentinel Physicians	Page 2
Children's Hospital ER	Page 2
Medical Services Plan	Page 3
Laboratory Surveillance	Page 5
ILI Outbreaks	Page 6
Pandemic H1N1 (pH1N1)	Page 7

Canada:

FluWatch Activity levels	Page 8
NML strain Characterization	Page 8
Anti-Viral Resistance	Page 8

International :

Northern Hemisphere	Page 8
Southern Hemisphere	Page 8

Other:

List of Acronyms	Page 9
Web Sites	Page 9
Outbreak Report Form	Page 10

Highlights

In week 37 (Sept 13-19), BC continued to experience an increase in influenza activity. The proportion of patients presenting to sentinel physicians with ILI and the proportion of Medical Services Plan claims for influenza illness increased compared to the previous week. Emergency room visits to BC Children's Hospital due to ILI remained approximately constant, at levels above those observed last year. One school outbreak and one long term care facility pH1N1 outbreak were reported during this period. At the BC Provincial Laboratory, 19.3% (49/374) of respiratory specimens were positive for influenza A, and all subtyped isolates were the pandemic H1N1 virus (pH1N1). Together surveillance indicators suggest that influenza activity is increasing and remains above the expected range for this time of year.

Report written: September 23, 2009

Edited: September 23, 2009

Disseminated/posted to web: Sept 23, 2009

Contributors: Vanita Sahni, Travis Hottes, Naveed Janjua, Danuta Skowronski

BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN


2008-09: Number 33, Week 37

September 13 – 19, 2009

British Columbia


Sentinel Physicians

During week 37, the percentage of patients presenting to sentinel physicians with ILI increased to 0.74%. This is substantially above the expected range for this time year. 74% (29/39) of sentinel physicians reported for week 37.



BC Children's Hospital Emergency Room

During week 37, 5.9% of Emergency Room visits to BC Children's hospital were attributed to ILI. This is similar to the previous week and above the proportion observed during the same time last year.



BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN


2008-09: Number 33, Week 37

September 13 – 19, 2009

Medical Services Plan

Influenza illness as a proportion of all submitted BC Medical Services Plan (MSP) claims increased in week 37. On a regional level, increases occurred in VIHA, VCH, and FHA. In these regions the proportion of claims for influenza is at or exceeding the 10 year maximum.

Influenza Illness Claims* British Columbia




*Influenza illness is tracked as the percentage of all submitted MSP general practitioner claims with ICD-9 code 487 (influenza).

**MSP week 27 Sep 2009 corresponds to sentinel ILI week 40.

***Current to September 15, 2009

Northern




BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN


2008-09: Number 33, Week 37

September 13 – 19, 2009


Interior




Vancouver Coastal



Fraser



Vancouver Island



BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN

2008-09: Number 33, Week 37


September 13 – 19, 2009

Laboratory Reports

There has been a large increase in the number of respiratory specimens submitted to BCCDC Laboratory Services. In week 37 they tested 590 respiratory specimens. 114 (19.3%) tested positive for influenza A (including pH1N1). Of those subtyped (n=102), 100% were pH1N1. No influenza B was detected. These proportions are similar to the previous week. Respiratory pathogens detected included: rhino/enterovirus (3% of specimens tested) and adenovirus (0.2%).

During week 37, Children's and Women's Health Centre Laboratory tested 72 respiratory specimens. Five were positive for influenza, 1 tested positive for parainfluenza and 1 for adenovirus.

Virus Detections and Percentage of Respiratory Specimens Submitted to BC Provincial Laboratory Diagnosed Positive for Influenza Virus, per Week, BC, 2008-2009




Note: The increase in bars during weeks 17-19 above reflects the large surge in specimens submitted to BCCDC for testing (2594 specimens were tested, a 5-fold increase over the number of tests performed during the 3-week period of peak activity this season).

BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN


2008-09: Number 33, Week 37

September 13 – 19, 2009



ILI Outbreaks

During week 37, one school outbreak was reported in VIHA, and a pH1N1 outbreak in a long term care facility was reported in VCH.



BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN

2008-09: Number 33, Week 37

September 13 – 19, 2009

Pandemic H1N1 (pH1N1)


BCCDC continues to monitor the pH1N1 virus pandemic. As of September 21, fifty-one cases in BC have been admitted to hospital. Among hospitalized cases, 63% had underlying medical conditions; 21% had lung disease, 15% had asthma and 10% had chronic heart disease. 39% (20) of hospitalized cases have been admitted to the intensive care unit and 12% (6) have died. As shown in the graph below, pH1N1 hospitalization rates are highest in those under 2 years of age.

For further description of BC pH1N1 cases, visit: www.bccdc.ca/dis-cond/DiseaseStatsReports/influSurveillanceReports.htm
 Resources for healthcare professionals: www.bccdc.ca/resourcematerials/newsandalerts/healthalerts/H1N1FluVirusHumanSwineFlu.htm


Pandemic H1N1*, BC Cases by Collection Date (as of September 21, 2009)

N = 964 (including 51 hospitalized cases)

* formerly known as swine-origin influenza virus



Cumulative Rate of pH1N1 Hospitalizations by Age, per 100,000 Population, BC April 17 - September 21, 2009



BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN

2008-09: Number 33, Week 37

September 13 – 19, 2009

CANADA

FluWatch

During week 36, national influenza activity levels increased slightly. Compared to week 35, ILI consultation rates increased from 14 to 23 consultations per 1000 patient visits; however this is within the expected range for this time of year. Compared to the previous week, the proportion of tests positive for influenza remained approximately constant at 2.5%, this represents an overall decline from 23% tests positive per patient visits in the week ending June 13. www.phac-aspc.gc.ca/fluwatch/

National Microbiology Laboratory

Since Sept 1, 2008 and as of August 31, 1306 influenza isolates from provincial and hospital labs have been characterized at the National Microbiology Laboratory (NML):

262 A/Brisbane/59/07(H1N1)-like* † from BC, AB, SK, MB, ON, QC, NB, NS, & PEI;

172 A/Brisbane/10/07(H3N2)-like* † from ten provinces;

11 B/Florida/04/06(Yamagata)-like* from AB, ON, QC & NB;

379 B/Malaysia/2506/04(Victoria)-like from all ten provinces;

180 B/ Brisbane/60/08(Victoria)-like † from BC, AB, SK, MB, ON, QC, NB, NS, & NU; and

302 A/California/07/2009-like[§] from BC, AB, SK, MB, ON, QC, NB, NS, NT, & NU;

* indicates a strain match to the 2008-09 vaccine

† indicates a strain match to the 2009-10 vaccine

[§] A/California/07/2009 (H1N1) is the variant reference virus (pH1N1) selected by WHO for a pandemic influenza A/H1N1 vaccine.

Antiviral Resistance

Drug susceptibility testing at the NML as of August 31 indicated that most (n=320) human influenza A/H1N1 isolates tested to date were resistant to oseltamivir (one human H1N1 isolate identified since mid-April was sensitive). All human H3N2 (n=194), influenza B (n=573), and pH1N1 (n=527) isolates tested at the NML were found to be sensitive to oseltamivir. Of the isolates tested for amantadine resistance, all (n=319) human H1N1 isolates were found to be sensitive, all (n=396) human H3N2 isolates were found to be resistant, and all (n=361) pH1N1 isolates were found to be resistant. All 1305 (257 human H1N1, 190 human H3N2, 578 influenza B, and 280 pH1N1) isolates that have been tested for zanamivir resistance were sensitive.

Global surveillance has shown that circulating pH1N1 viruses are resistant to amantadine but remain sensitive to zanamivir and oseltamivir, although sporadic cases of oseltamivir resistance have been observed worldwide. The first cases of oseltamivir resistance with an epidemiological link were identified in the US on August 14 and 19.

INTERNATIONAL

Northern Hemisphere: In the United States, in the week ending September 12 influenza activity as determined by sentinel physician visits and geographic spread increased. 4.4% of visits to sentinel physicians were for ILI, an increase from 1.4% in week 33. Eighteen percent of respiratory specimens tested in reference laboratories during this week were positive for influenza, this proportion was similar to the previous week. Ninety-nine percent of the subtyped influenza A viruses were pH1N1. In Europe for the week ending September 13, influenza activity remains low or declining in most countries, with the exception of Sweden where there is medium, widespread activity with an increasing trend. <http://www.cdc.gov/flu/weekly/> and <http://www.eiss.org>.

Southern Hemisphere: Several countries in the Southern Hemisphere previously reporting severe winter influenza activity have now passed the peak. Notably as of September 4th in Australia, influenza activity is continuing to decrease and presentations to sentinel physicians and ERs is below the previous years average. In New Zealand as of September 13th, pH1N1 activity continues to decline; consultations with sentinel physicians have declined to less than a third of those observed during the peak in early July, and are now approaching baseline levels. In Chile as of September 9, ILI activity is within the range expected for this time of year. In South Africa, laboratories are currently reporting pH1N1 as the dominant influenza subtype; previously, in June and July of this year the dominant subtype was A/H3N2.

BRITISH COLUMBIA INFLUENZA SURVEILLANCE BULLETIN

2008-09: Number 33, Week 37

September 13 – 19, 2009

Contact Us:

Epidemiology Services : BC Centre for Disease Control (BCCDC)

655 W. 12th Ave, Vancouver BC V5Z 4R4. Tel: (604) 707-2510 / Fax: (604) 707-2516. InfluenzaFieldEpi@bccdc.ca

List of Acronyms

ACF: Acute Care Facility

AI: Avian Influenza

FHA: Fraser Health Authority

HMPV: Human metapneumovirus

HSDA: Health Service Delivery Area

IHA: Interior Health Authority

ILI: Influenza-Like Illness

LTCF: Long Term Care Facility

MSP: BC Medical Services Plan

NHA: Northern Health Authority

NML: National Microbiological Laboratory

pH1N1: Pandemic H1N1 influenza or swine origin influenza

RSV: Respiratory syncytial virus

VCHA: Vancouver Coastal Health Authority

VIHA: Vancouver Island Health Authority

WHO: World Health Organization

Web Sites

1. Influenza Web Sites

Canada – Flu Watch: www.phac-aspc.gc.ca/fluwatch/

Washington State Flu Updates: www.doh.wa.gov/ehsphl/epidemiology/CD/HTML/FluUpdate.htm

USA Weekly Surveillance reports: www.cdc.gov/flu/weekly/

European Influenza Surveillance Scheme: www.eiss.org/index.cgi

WHO – Global Influenza Programme: www.who.int/csr/disease/influenza/mission/

WHO – Weekly Epidemiological Record: www.who.int/wer/en/

Influenza Centre (Australia): www.influenzacentre.org/

Australian Influenza Report: www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm

New Zealand Influenza Surveillance Reports: www.surv.esr.cri.nz/virology/influenza_weekly_update.php

2. Avian Influenza Web Sites

World Health Organization – Avian Influenza: www.who.int/csr/disease/avian_influenza/en/

World Organization for Animal Health: www.oie.int/eng/en_index.htm

3. Pandemic H1N1 Influenza Web Sites

BCCDC: www.bccdc.ca/dis-cond/a-z/_h/HumanSwineFlu/default.htm

PHAC: www.phac-aspc.gc.ca/alert-alerte/swine_200904-eng.php

US CDC: www.cdc.gov/swineflu/index.htm

WHO: www.who.int/csr/disease/swineflu/en/index.html

4. This Report On-line: www.bccdc.ca/dis-cond/DiseaseStatsReports/influSurveillanceReports.htm

Influenza-Like Illness (ILI) Outbreak Summary Report Form

Please complete and email to ilioutbreak@bccdc.ca or fax to (604) 707-2516

ILI: Acute onset of respiratory illness with fever and cough and with one or more of the following: sore throat, arthralgia, myalgia, or prostration which *could* be due to influenza virus. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

Schools and work site outbreak: greater than 10% absenteeism on any day, most likely due to ILI.

Residential institutions (facilities) outbreak: two or more cases of ILI within a seven-day period.

SECTION A: Reporting Information

Person Reporting: _____ Title: _____

Contact Phone: _____ Email: _____

Health Authority: _____ HSDA: _____

Full Facility Name: _____

- Is this report: First Notification (*complete section B below; Section D if available*)
 Update (*complete section C below; Section D if available*)
 Outbreak Over (*complete section C below; Section D if available*)

SECTION B: First Notification

Type of facility: LTCF Acute Care Hospital Senior's Residence
(if ward or wing, please specify name/number: _____)
 Workplace School (grades: _____) Other (_____)

Date of onset of first case of ILI (dd/mm/yyyy): _____ / _____ / _____

Numbers to date	Residents/Students	Staff
Total		
With ILI		
Hospitalized		
Died		

SECTION C: Update AND Outbreak Declared Over

Date of onset for most recent case of ILI (dd/mm/yyyy): _____ / _____ / _____

If over, date outbreak declared over (dd/mm/yyyy): _____ / _____ / _____

Numbers to date	Residents/Students	Staff
Total		
With ILI		
Hospitalized		
Died		

SECTION D: Laboratory Information

Specimen(s) submitted? Yes (location: _____) No Don't know
 If yes, organism identified? Yes (specify: _____) No Don't know