



BC Centre for Disease Control
An agency of the Provincial Health Services Authority

655 West 12th Avenue
Vancouver, BC V5Z 4R4

Tel 604.707.2443
Fax 604.707.2441

www.bccdc.ca

Clinical Prevention Services
Provincial STI Services

604.707.5600
604.707.5604

www.SmartSexResource.com

Ceftriaxone

Ceftriaxone is an antibiotic given by injection into a muscle for the treatment of certain sexually transmitted infections (STI).

Allergies

- Tell your healthcare provider if you have an allergy to ceftriaxone, other cephalosporin antibiotics such as cefixime (Suprax®), cephalexin (Keflex®), cefaclor (Ceclor®) or penicillin antibiotics such as penicillin V-K, amoxicillin (Amoxil®) or to lidocaine, or other local anesthetics or drugs of the amide type (e.g., prilocaine, mexiletine, amiodarone). Lidocaine is mixed with ceftriaxone to reduce pain at the injection site.

Pregnancy and Chest/Breastfeeding

Ceftriaxone may be used with caution during pregnancy and chest/breastfeeding. Please consult your healthcare provider.

CAUTION

- **Tell your healthcare provider if you have any gallbladder, liver, kidney, pancreas or bleeding disorder, or a history of stomach dysfunction, particularly colitis.**
- **Do not take the following medications with ceftriaxone:**
 - Oral Typhoid vaccine (Vivotif®)
- **Ceftriaxone may interfere with certain urine glucose tests.** (Clinitest®)
- **Ceftriaxone is contraindicated in preterm newborns or infants less than one month old with high blood levels of bilirubin.**
- **Drug Interactions:** Please note that individual drug interactions are no longer listed in this document. If you are taking any prescription, non-prescription, herbal, or recreational products, please discuss with your healthcare provider.

Side Effects

- You may get discomfort, redness or swelling at the injection site.
- You may experience diarrhea.
- If any of these effects persist or worsen, please contact your healthcare provider.

Ceftriaxone

Care Following Treatment

- Please wait in the clinic 15 minutes after your injection.
- If you have any reaction following treatment (e.g., rash, itchiness, difficulty breathing), please inform a staff member immediately.
- If these symptoms occur after leaving the clinic, please seek emergency medical help immediately.

Special Instructions

- Do not have sex until:
 - One week after you complete your treatment with ceftriaxone , or until
 - Your sex partner(s) have also been treated, even if their test results are negative.
- You will need retreatment if you have sex with an untreated partner, or if you have sex before either you or your partner's treatment is complete. Please discuss with your healthcare provider.

If you have any questions or need further information, please visit www.smartsexresource.com or contact your healthcare provider.