

# BC ROUTINE IMMUNIZATION SCHEDULE INFANTS & CHILDREN

VACCINE	CHILD'S AGE					
	2 months	4 months	6 months	12 months	18 months	Starting at 4 years (kindergarten entry)
<b>DTaP-HB-IPV-Hib</b> (diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # <a href="#">105</a> 2. Product Monograph <a href="#">Infanrix Hexa</a>	✓	✓	✓			
<b>Pneumococcal conjugate ‡</b> 1. HealthLinkBC File # <a href="#">62a</a> 2. Product Monograph <a href="#">Prennar 13</a>	✓	✓		✓		
<b>Rotavirus</b> 1. HealthLinkBC File # <a href="#">104a</a> 2. Product Monograph <a href="#">Rotarix</a>	✓	✓				
<b>Meningococcal C conjugate</b> 1. HealthLinkBC File # <a href="#">23a</a> 2. Product Monograph <a href="#">NeisVac-C</a>	✓			✓		
<b>MMR</b> (measles, mumps, rubella) 1. HealthLinkBC File # <a href="#">14a</a> 2. Product Monograph <a href="#">MMR II</a> / <a href="#">Priorix</a>				✓		
<b>Varicella †</b> (chickenpox) 1. HealthLinkBC File # <a href="#">44b</a> 2. Product Monograph <a href="#">Varilrix</a> / <a href="#">Varivax III</a>				✓		
<b>DTaP-IPV-Hib</b> (diphtheria, tetanus, pertussis, polio, <i>Haemophilus influenzae</i> type b) 1. HealthLinkBC File # <a href="#">15b</a> 2. Product Monograph <a href="#">Infanrix-IPV/Hib</a> / <a href="#">PediaceL</a>					✓	
<b>Tdap-IPV</b> (tetanus, diphtheria, pertussis, polio) 1. HealthLinkBC File # <a href="#">15a</a> 2. Product Monograph <a href="#">Adacel-Polio</a> / <a href="#">Boostrix-Polio</a>						✓
<b>MMRV †</b> (measles, mumps, rubella, varicella) 1. HealthLinkBC File # <a href="#">14e</a> 2. Product Monograph <a href="#">Priorix-Tetra</a> / <a href="#">ProQuad</a>						✓
<b>Influenza *</b> 1. HealthLinkBC File # <a href="#">12d</a> and <a href="#">12e</a> 2. Product Monograph <a href="#">FluLaval Tetra</a> / <a href="#">Flumist Quadrivalent</a> / <a href="#">Fluzone Quadrivalent</a>			✓ (yearly for children 6 months of age and older)			
<b>Hepatitis A</b> (for Indigenous children only) 1. HealthLinkBC File # <a href="#">33</a> 2. Product Monograph <a href="#">Havrix</a> / <a href="#">Vaqta</a>			✓		✓	

‡ Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. For more information see the BC Immunization Manual, [Part 4 – Biological Products](#).

† Children who had chickenpox or shingles disease, confirmed by a lab test, at 1 year of age or older do not need the chickenpox vaccine and should get the MMR vaccine at 4 years of age.

\* A second dose is needed 4 weeks after the first dose if receiving the vaccine for the first time.

Access the [BC Immunization Manual](#)